

# Tattooing and body piercing guidance

---

# Toolkit

---



Chartered  
Institute of  
Environmental  
Health



## Tattooing and body piercing guidance

### ACKNOWLEDGEMENTS

Click on text to view

[FOREWORD](#)

[ENDORISING ORGANISATIONS](#)

[INTRODUCTION](#)

[USING THE GUIDANCE  
TOOLKIT](#)

[ACKNOWLEDGEMENTS](#)

#### **Acknowledgements**

These guidelines are extensively based on the Tattooing and Body Piercing Guidance: Toolkit which was published by the Chartered Institute of Environmental Health (CIEH) London in 2013. We acknowledge the contributions of the original authors and also those who contributed to the production of this Northern Ireland version as set out below:

---

#### **TATTOOING AND SKIN PIERCING WORKING GROUP MEMBERS**

---

**Chartered Institute of Environmental Health**

**Department of Health, Social Services and Public Safety (DHSSPS)**

**Public Health Agency**

**Health and Safety Liaison Group for Northern Ireland**

**Subgroup of the Chief Environmental Health Officers Group  
(CEHOG)**

January 2014

# Tattooing and body piercing guidance

<b>FOREWORD</b>	<b>2</b>	<b>Appendix 03</b>	<b>42</b>
<b>ENDORISING ORGANISATIONS</b>	<b>3</b>	Safe use and disposal of sharps	
<b>INTRODUCTION</b>	<b>4 – 5</b>	<b>Appendix 04</b>	<b>43</b>
<b>USING THE GUIDANCE TOOLKIT</b>	<b>6</b>	First Aid following a blood/body fluid exposure	
<b>PART A</b>			
<b>Section 01</b>	<b>7 – 12</b>	<b>Appendix 05</b>	<b>44</b>
Legislative background on tattooing and skin piercing activities		Protocol for cleaning up blood or a blood stained body fluid spill	
<b>Section 02a</b>	<b>13</b>	<b>Appendix 06</b>	<b>45</b>
Standard Principles of Infection Control		Principles for good waste handling	
<b>Section 02b</b>	<b>14 – 15</b>	<b>Appendix 07</b>	<b>45</b>
Principles of Infection Control – Hand hygiene		Template protocol for environmental cleaning of premises	
<b>Section 02c</b>	<b>16 – 18</b>	<b>Appendix 08</b>	<b>46</b>
Principles of Infection Control – Personal protective equipment		Tattooing/body piercing consent form	
<b>Section 02d</b>	<b>19 – 21</b>	<b>Appendix 09</b>	<b>48</b>
Principles of Infection Control – Management of sharps and exposure to blood and body fluids		Aftercare follow-up record sheet	
<b>Section 02e</b>	<b>22 – 23</b>	<b>Appendix 10</b>	<b>49</b>
Principles of Infection Control – Safe handling, storage and disposal of waste materials		Decontamination requirements for equipment used in tattooing and skin piercing	
<b>Section 02f</b>	<b>24</b>	<b>Appendix 11</b>	<b>50</b>
Principles of Infection Control – Cleaning and disinfection of the environment		Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners	
<b>Section 02g</b>	<b>25</b>	<b>Appendix 12</b>	<b>51</b>
References		Equipment and body piercing jewellery sterilization standard for tattooists and body piercers	
<b>Section 03</b>	<b>26 – 27</b>	<b>Appendix 13</b>	<b>52</b>
Before and aftercare of a tattoo or body piercing		Autoclave daily record sheet	
<b>Section 04</b>	<b>28 – 32</b>	<b>PART C</b>	
Decontamination		<b>Leaflets to download and print out</b>	
<b>Section 05</b>	<b>33 – 34</b>	01 Tattoo aftercare	53
Product quality of tattoo ink		02 Ear and face piercing aftercare	54
<b>Section 06</b>	<b>35</b>	03 Oral piercing aftercare	55
Body piercing jewellery		04 Body and surface piercing aftercare	56
<b>Section 07</b>	<b>36 – 37</b>	05 Genital piercing (female) aftercare	57
Governance		06 Genital piercing (male) aftercare	58
<b>Section 08</b>	<b>38</b>	07 Microdermal implants aftercare	59
Management of infectious disease incidents relating to tattooing and body piercing		<b>Poster to download</b>	
<b>PART B</b>			
<b>Appendix 01</b>	<b>40 – 41</b>	How to handwash	60
Infection, its causes and spread, including a glossary of infection-related terms		<b>PART D</b>	
<b>Appendix 02</b>	<b>41</b>	<b>Audit Tool to download</b>	
Blood borne viruses		<b>61</b>	
<b>PART E</b>			
<b>Literature review to download</b>			
<b>62</b>			

# Tattooing and body piercing guidance

## FOREWORD

### **GARY MC FARLANE**

**Director, Chartered Institute of  
Environmental Health Northern Ireland**

Tattooing and body piercing have become increasingly popular and fashionable. Ensuring practitioners follow safe working practices is important for protection of both clients and the practitioners themselves. This toolkit has been developed specifically for Northern Ireland following the publication of a similar resource for England. In doing so we have collaborated and worked with the relevant NI agencies, notably the Department of Health Social services and Public Safety (DHSSPS), the Public Health Agency and the Health and Safety Liaison Group (HSLG), a subgroup of the Chief Environmental Health Officers Group (CEHOG). We hope that the resources will prove a practical tool for practitioners.

### **PATRICIA ALLEN**

**Assistant Director of Northern Group  
Systems Environmental Health**

This guidance toolkit aims to influence tattooing and body piercing practices in Northern Ireland, in particular by the promotion of evidence based prevention and control measures. It provides easy access to key guidance and best practice, giving practitioners and others a single authoritative source of information. By doing so it is hoped that standards of safety and hygiene delivered across the range of tattooing and body piercing practices will be consistently high for Northern Ireland consumers.

### **DR LORRAINE DOHERTY**

**Assistant Director for Health Protection  
Public Health Agency**

The tattooing and body piercing industry has seen a period of growth in Northern Ireland in recent years. As such there is now an increased need to ensure that those who are working in this field are supported in delivering safe practice. As an interactive document, this toolkit acts as a framework for good infection and control practice which enables the user to access the sound evidence based guidance and to appropriately manage infection risk. I would like to acknowledge the excellent collaborative working of health protection colleagues from PHA, PHE and CIEH and colleagues from the tattooing industry that has resulted in the publication of this invaluable tool. It is my pleasure to endorse this guidance toolkit as a key resource for practitioners in the industry which can only lead to better health outcomes for service users.

### **NIGEL MC MAHON**

**Chief Environmental Health Officer  
Department of Health, Social Services &  
Public Safety**

Infection prevention and control is a paramount public health concern. The promotion of safe working practices is important for practitioners, clients and public health professionals including environmental health staff. This reference guide attempts to bring all of the relevant information together in one place for the benefit of those involved in the industry, as well as for those that seek to advise and regulate it. The guide has been produced in partnership with a number of organisations and individuals and I would offer my thanks to all those involved. On behalf of the working group, I would also like to record our appreciation of the Chartered Institute of Environmental Health for carrying out the editing, web design and publication of the toolkit online.

# Tattooing and body piercing guidance

## ENDORISING ORGANISATIONS

Click on text to view

[FOREWORD](#)

[ENDORISING ORGANISATIONS](#)

[INTRODUCTION](#)

[USING THE GUIDANCE TOOLKIT](#)

### **CHARTERED INSTITUTE OF ENVIRONMENTAL HEALTH (CIEH)**

The CIEH is a registered charity and the professional voice for environmental health. It sets standards, accredits courses and qualifications for the education of members and other environmental health practitioners. It provides information, evidence and policy advice to local and national government and environmental and public health practitioners in the public and private sectors. As an awarding body, the CIEH provides qualifications, events, and support materials on topics relevant to health, wellbeing and safety to develop workplace skills and best practice.

### **THE DEPARTMENT OF HEALTH, SOCIAL SERVICES AND PUBLIC SAFETY (DHSSPS)**

DHSSPS was established by the Departments (NI) Order 1999. It is the Department's mission to improve the health and social well-being of the people of Northern Ireland.

The Department has three main business responsibilities:

- **Health and Social Care (HSC)**, which includes policy and legislation for hospitals, family practitioner services and community health and personal social services;
- **Public Health** which covers policy, legislation and administrative action to promote and protect the health and well-being of the population; and
- **Public Safety**, which covers policy and legislation for fire and rescue services.

### **THE PUBLIC HEALTH AGENCY (PHA)**

From its establishment in 2009, Public Health Agency provides a renewed and enhanced focus on public health and wellbeing by bringing together a wide range of public health functions under one organisation.

PHA is a multi-disciplinary, multi-professional body with a strong regional and local presence. It has four key functions:

- health and social wellbeing improvement;
- health protection;
- public health support to commissioning and policy development ;
- HSC research and development

The PHA also work to create better inter-sectoral working, including enhanced partnership arrangements with local government, to tackle the underlying cause of poor health and reduce health inequalities.

### **NORTHERN IRELAND CHIEF ENVIRONMENTAL HEALTH OFFICERS GROUP**

The Northern Ireland Chief Environmental Officers Group (CEHOG) was formed as a liaison body to provide a forum to unite the local government environmental health service in Northern Ireland and to establish and maintain effective services partnerships with bodies having an influence upon health in NI communities. The fundamental remit is to aid the co-ordination and consistency of environmental health services, to assist the development of highly quality services and to provide a consultative body with links to other agencies and departments to facilitate the passage of advice on relevant policy matters.

Membership of this liaison body includes all twenty six District Chief Environmental Health Officers, (or equivalent in title and function), the four Group Chief Environmental Health Officers and from outside local government, the Chief Environmental Health Officer, DHSSPS and the NI Director of the Chartered Institute of Environmental Health.

# Tattooing and body piercing guidance

## INTRODUCTION

Click on text to view

[FOREWORD](#)

[ENDORISING ORGANISATIONS](#)

**[INTRODUCTION](#)**

[USING THE GUIDANCE TOOLKIT](#)

This guidance toolkit has been prepared by a panel of health protection and practitioner representatives. It comprises a consensus of expert advice which it is intended will provide an authoritative source of information. Its contents are supported by extensive literature reviews (Part E).

The purpose of the guidance toolkit is to support local authority and other regulatory officers in determining their requirements for effective control of risk in these activities and to promote a consistent approach. Similarly, it is intended to be of assistance to practitioners and businesses undertaking these activities to support them in adopting acceptable standards of practice. The use of this guidance toolkit will help to ensure the health and safety of both clients and operators and that tattooing and skin piercing practitioners will be operating in compliance with legal requirements.

### BACKGROUND

Tattooing and body piercing procedures have become more popular and fashionable in the United Kingdom (UK) as a whole, particularly in the last decade. The range of tattooing and body piercing procedures has also increased. There are no published data on the prevalence of tattoos in the general UK population. There are also no comprehensive data for the UK on the prevalence of body piercing, but a small study has estimated that the prevalence of body piercing, other than of earlobes, in the general adult population in England was 10% (Bone A et al, 2008).

There are known and well reported health risks which can be attributed to these procedures, as well as associated legal issues. Improper and unhygienic practice may result in localised skin infections at the site of the tattoo or piercing. There is also the risk of transmission of blood-borne viruses, for example Hepatitis B, Hepatitis C, Hepatitis D or HIV, which can have more serious and long term health consequences. It is therefore important that practitioners have safe working practices, and particularly that good infection control practices are followed at all times, so that both clients and practitioners are protected.

Over the years a variety of legislation has been introduced, mainly for local adoption, to encourage and support safe practice. Model byelaws have been made available, but have not been accompanied by standard requirements for compliance. As a response to on-going concerns, a number of sets of local/regional guidelines have been developed by different agencies, often initiated by environmental health or health protection specialists (this guidance toolkit draws upon some of the previously published material). However, there have been difficulties previously in engaging practitioners in the development of such guidelines and in securing adoption and wider implementation.

It is also recognised that there are no nationally recognised or accredited training courses, standards for practice, agreed knowledge and skills frameworks or arrangements for monitoring and reporting of professional competence. The absence of accredited training and competencies for tattooing and body piercing is an area that needs to be addressed nationally and is outside the scope of this guidance. The working group has, however, been greatly assisted in the preparation of this guidance by the Tattoo and Piercing Industry Union who are recognised as a professional body for tattoo and body piercing practitioners in the UK.



## Tattooing and body piercing guidance

### INTRODUCTION (*continued*)

Click on text to view

[FOREWORD](#)

[ENDORISING ORGANISATIONS](#)

[INTRODUCTION](#)

[USING THE GUIDANCE  
TOOLKIT](#)

#### **DEVELOPMENT OF THE GUIDANCE FOR NORTHERN IRELAND**

This guidance was developed in response to concerns raised by tattoo and body piercing practitioners, as well as health protection and environmental health specialists. These concerns were particularly in regard to the lack of robust and consistent guidance on standards of hygiene and safety. This has been leading to inconsistency in advice and variations in standards of practice.

A multi-agency steering group was set up comprising representatives from the Chartered Institute of Environmental Health, Department of Health and Social Services and Public Safety (DHSSPS), Public Health Agency (PHA) and the Health and Safety Liaison Group for Northern Ireland (a subgroup of the Chief Environmental Health Officers Group) and also individuals with practical experience of working in this area as expert advisors, practitioners or regulators.

The guidance is supported by extensive documentary evidence of scientific knowledge, reported research and published literature encompassing expert advice and the opinions and experience of practitioners of what works at a practical level.

#### **References**

Bone A., Ncube F., Nichols T. & Noah ND. (2008) Body piercing in England: a survey of piercing at sites other than earlobe. *BMJ*; 336; 1426-1428.

# Tattooing and body piercing guidance

## USING THE GUIDANCE TOOLKIT

Click on text to view

[FOREWORD](#)

[ENDORISING ORGANISATIONS](#)

[INTRODUCTION](#)

[USING THE GUIDANCE  
TOOLKIT](#)

[THE AUDIT TOOL](#)

[PROVIDING FEEDBACK](#)

The guidance has been written as a key point of reference for use nationally by tattoo and body piercing practitioners who work in regulated premises, local authority officers in their regulatory role and health protection staff who are asked to provide expert advice. The guidance does not cover mobile operators or non-registered practitioners, although the risks encountered in relation to their activities will be of equal or greater concern. It does not address the risks associated with procedures other than those commonly accepted as necessary for tattooing and the insertion of body jewellery, although the procedures recommended for infection control are based upon sound principles of infection control and will have wider application.

The material is arranged so as to be readily accessible as a web-based toolkit, organised in a manner that reflects the tattoo and body piercing setting, and with supportive documentation and literature that can be downloaded and saved or printed in the manner that users prefer.

It is intended that the adoption of the standards recommended in this guidance, particularly those relating to infection control and decontamination, will help to establish standards for good practice. Governance is promoted by the inclusion of template consent forms, aftercare advice leaflets and a good practice infection control audit tool.

### THE AUDIT TOOL

This audit tool can be downloaded and used as individual sections or as a complete tool. It is intended to be used by tattoo and body piercing practitioners, regulatory officers and health protection practitioners to generate evidence of the environment, practice and procedures in meeting standards and whether practitioners are applying best practice and following guidance. The tool can be repeated to see if standards have been maintained or improved.

No copyright is being claimed for the toolkit or any of the material it contains and the authors encourage its wider distribution and use.

In offering and using the advice contained in this guidance it must be clearly understood that:

- Legislation may change over time and the advice given is based on the information available at the time this toolkit was produced – it is not necessarily comprehensive and is subject to revision in the light of further information.
- Only the courts can interpret legislation with any authority, and
- This advice is not intended to be definitive legal guidance nor is it a substitute for the relevant law and independent legal advice should be sought where appropriate.

### PROVIDING FEEDBACK

We hope you find the guidance useful. It is intended that the document will be revisited and updated periodically. To that end feedback on your experience is positively encouraged. Feedback can be sent to Gary Mc Farlane, [g.mcfarlane@cieh.org](mailto:g.mcfarlane@cieh.org) Director of The Chartered Institute of Environmental Health Northern Ireland.



Click on text to view

#### SUMMARY

[Specific controls](#)

[General controls](#)

[Determining requirements](#)

#### REGISTRATION AND BYELAW REQUIREMENTS IN NORTHERN IRELAND

#### HEALTH AND SAFETY AT WORK (NI) ORDER 1978

[General duties](#)

[Risk assessment](#)

[Control of substances hazardous to health](#)

[Management of contractors](#)

[Enforcement](#)

#### AGE LIMITS AND CONSENT

[Tattooing](#)

[Other skin piercing activities](#)

[Acupuncture and electrolysis](#)

[Consent](#)

#### USE OF LOCAL ANAESTHETIC MEDICATION

# Tattooing and body piercing guidance

## PART A

### Legislative background on tattooing and skin piercing activities

#### SUMMARY

The use of legislation in this area of activity is primarily to ensure that infection control arrangements are adequate and effectively carried out wherever tattooing and skin piercing are carried out.

The primary means of enforcing infection control arrangements is by use of registration and the observance of provisions contained in byelaws. The registration and byelaw provisions are largely concerned with setting requirements for good standards by requiring the maintenance of established hygiene controls in respect of premises, equipment, procedures and practices. However, there are additional controls contained in primary legislation that do contain provisions for the immediate prohibition of activities or persons or for the closure of premises where risk of infection can be demonstrated.

The legislation relating to tattooing and skin piercing activities, can therefore be broadly split into two main areas:

- Specific controls by registration of premises and people carrying out the activities, and associated byelaws, and
- General controls of activities through primary legislation that is not specific to particular activities but applies to all of them.

#### Specific controls

Arrangements for registration will differ depending on the particular requirements of the local authority in whose area the business is located or the activity is being carried out. The majority of local authorities have adopted byelaws in respect of the standards for the maintenance of established hygiene controls in respect of premises, equipment, procedures and practices.

#### General controls

The primary health and safety legislation in Northern Ireland (NI) is the Health and Safety at Work (Northern Ireland) Order 1978 (HSWO). It can be used to impose and enforce infection control requirements in relation to all skin piercing activities, including peripatetic practitioners who visit a client's home. It allows for immediate prohibition of persons or activities that pose an imminent risk to health or safety.

#### Determining requirements

It is intended that this guidance will provide an additional authoritative source of information to support local authorities in determining their requirements for effective control of risk in these activities and adopting a consistent approach in the application of the legislation. Similarly, it will be of assistance to those businesses and practitioners undertaking these activities to ensure that they are able to operate safely and comply with legal requirements.

Click on text to view

**SUMMARY**

Specific controls

General controls

Determining requirements

**REGISTRATION AND BYELAW REQUIREMENTS IN NORTHERN IRELAND****HEALTH AND SAFETY AT WORK (NI) ORDER 1978**

General duties

Risk assessment

Control of substances  
hazardous to health

Management of contractors

Enforcement

**AGE LIMITS AND CONSENT**

Tattooing

Other skin piercing activities

Acupuncture and electrolysis

Consent

**USE OF LOCAL ANAESTHETIC MEDICATION**

# Tattooing and body piercing guidance

## PART A

### Legislative background on tattooing and skin piercing activities (*continued*)

#### REGISTRATION AND BYELAW REQUIREMENTS IN NORTHERN IRELAND

There are provisions in Part V of the [The Local Government \(Miscellaneous Provisions\) Northern Ireland Order 1985](#) (LGMPO) for local authorities in NI to require the registration of persons carrying on the practices of acupuncture, tattooing, ear piercing or electrolysis. These powers are adoptive, and local authorities are able to choose which of these practices would be required to be registered in their area. [The Local Government \(Northern Ireland\) Order 2005](#) added semi-permanent skin-colouring and cosmetic piercing to this list of activities for which registration can be required.

The Order allows for local authorities to make byelaws, for the purpose of securing; the cleanliness and hygiene of premises, practitioners and equipment. These measures are intended to increase health protection and reduce the risk of transmission of blood borne virus infections.

The local authority can request reasonable information from applicants for registration. This cannot include details concerning persons whom the applicant has given treatments to, however, it could include evidence of training or competency for those being registered.

The registration is mandatory in that the local authority must issue a registration if the application has been properly made. A registration can only be refused where a person has previously been convicted of an offence under Article 15 of the 1985 Order and the convicting magistrate suspended or cancelled the previous registration. A registration can only be cancelled by a magistrate upon conviction of an offence, and this is instead of or in addition to a fine.

Article 15 of the 1985 Order provides for offences and for non-custodial penalties (summary conviction and fine) for trading without council and registration or breaching council byelaws. The court may also order suspension of, or cancellation of registration (whether of a person or premises) on conviction. When cancellation of registration happens, the court may order a fine, increased on a daily basis for late surrender of the cancelled registration certificate. There is also an offence of not displaying a certificate of registration or byelaws (in respect of which a person is liable on summary conviction to a fine).

The penalties for offences are fines on the Standard Scale of Level 3 (currently £1000) for offences under Article 15(1), Article 15(2), and Level 1 (currently £200) for an offence under Article 15(9).

There are some exemptions from the registration requirements. It does not apply to practices carried out by or under the supervision of a person who is registered as a medical practitioner (a Doctor registered with the General Medical Council) or for acupuncture by a dentist, or chartered physiotherapist, or a state registered physiotherapist, or a state registered chiroprapist, or to premises under their supervision.

Local authority officers can be authorised to enter any premises where they have reason to believe that an offence under Article 15 is being committed there, but where entry is refused the authority of a warrant issued by a Justice of the Peace has to be obtained.

The Department of Health Social Services and Public Safety (DHSSPS) issued guidance on the legal provisions relating to the regulation of cosmetic piercing and skin colouring businesses in 2005. The document also provides model byelaws which local authorities could formally adopt as well as guidance on the procedure for the confirmation of byelaws by the DHSSPS.

<http://www.dhsspsni.gov.uk/bodypiercing-order.pdf>

## Section 01

Click on text to view

**SUMMARY****Specific controls****General controls****Determining requirements****REGISTRATION AND BYELAW REQUIREMENTS IN NORTHERN IRELAND****HEALTH AND SAFETY AT WORK (NI) ORDER 1978****General duties****Risk assessment****Control of substances****hazardous to health****Management of contractors****Enforcement****AGE LIMITS AND CONSENT****Tattooing****Other skin piercing activities**

Acupuncture and electrolysis

**Consent****USE OF LOCAL ANAESTHETIC MEDICATION**

# Tattooing and body piercing guidance

## PART A

### Legislative background on tattooing and skin piercing activities (*continued*)

#### **HEALTH AND SAFETY AT WORK (NI) ORDER 1978**

The [Health & Safety at Work \(Northern Ireland\) Order 1978](#) (HSWO) applies to all persons engaged in tattooing and skin piercing activities for gain or reward. This includes peripatetic workers who carry out treatments in the client's home, although only the Health & Safety Executive for Northern Ireland have powers in relation to peripatetic workers.

It provides means of securing effective infection control and the following areas are particularly applicable.

#### **General duties**

Under Article 4 of the Order, all employers have a general duty of care to ensure the health, safety and welfare of their employees. Under Article 5, both employers and self-employed persons have a general duty of care to ensure their activities do not expose them or the general public to risks to their health or safety.

#### **Risk assessment**

A risk assessment is the key step in protecting workers and the public, as well as complying with the law. The risk assessment is a careful examination of what work activities could cause harm to people and this then guides decisions about precautions that need to be taken, including infection prevention and control measures.

The HSE provides detailed advice on carrying out risk assessments <http://www.hse.gov.uk/risk/risk-assessment.htm>, including interactive tools <http://www.hse.gov.uk/risk/shop.htm> and other pieces of guidance <http://www.hseni.gov.uk/guidance/content-getting-started/content-risk-assessment.htm>.

The [Management of Health and Safety at Work Regulations \(Northern Ireland\) 2000](#) requires all employers and self employed persons to:

- Undertake a risk assessment of their activities;
- Remove, where possible, that risk or;
- Where residual risk is unavoidable, to provide control measures to reduce it as far as possible, including as a last resort, provision of personal protective equipment;
- Provide training to staff and persons they use to undertake their business activities (contractors) to ensure they understand the risks and the control measures.

One of the risks that must be considered is that of possible complications relating to skin piercing and tattooing. The practitioner must make sure that a fully 'informed consent' procedure is adopted. This means gathering information from the client about their health and suitability for the treatment, and giving the client enough information about the possible complications that could arise from the treatment for them to make their own decision.

#### **Control of substances hazardous to health**

The [Control of Substances Hazardous to Health Regulations \(Northern Ireland\) 2003 \(COSHH\)](#) requires that a specific risk assessment is carried out by employers or self employed persons who work with substances hazardous to health. Substances which are hazardous to health include biological agents. The hazards in this context are the organisms which can cause communicable diseases could be transmitted from person to person by unhygienic practices.

Therefore a specific risk assessment in respect of infection control is necessary for all persons undertaking tattooing and skin piercing activities.

Businesses employing less than 5 people do not have to record the findings of this risk assessment, however they still have to satisfy regulatory officers that their risk assessment is suitable and sufficient.

Click on text to view

**SUMMARY****Specific controls****General controls****Determining requirements****REGISTRATION AND****BYELAW REQUIREMENTS IN NORTHERN IRELAND****HEALTH AND SAFETY AT WORK (NI) ORDER 1978****General duties****Risk assessment****Control of substances****hazardous to health****Management of contractors****Enforcement****AGE LIMITS AND CONSENT****Tattooing****Other skin piercing activities**

Acupuncture and electrolysis

**Consent****USE OF LOCAL ANAESTHETIC MEDICATION**

# Tattooing and body piercing guidance

## PART A

### Legislative background on tattooing and skin piercing activities (*continued*)

#### Management of contractors

The [Management of Health and Safety at Work Regulations \(Northern Ireland\) 2000](#) (Regulations Sections 11, 12 & 13) contains legal provisions which can be of particular importance to the many tattooists and cosmetic piercers who do not own the premises which they operate from and are not employed by the person who owns and/or manages the premises where they work. In these cases, they are usually self-employed and have some form of contract, formal or informal, with the owner of the shop premises whereby they pay the owner to allow them to work there. In these circumstances the shop owner can be said to be contracting out to the practitioner and in effect 'endorsing' their work'.

These tattooists and piercers are therefore contractors undertaking the activities of the business owner for them. This means that the premises owners cannot abdicate their own general duties under the HSWO to the individual practitioners. They have a duty to ensure that persons working on their premises are competent and that they carry out their work in a safe manner. The only way they can do this satisfactorily is to assess the practitioners for themselves and monitor their activities to ensure they have carried out their own risk assessment, as they are required to do by law, and that they are following control measures they have identified. The business owner has the ultimate power to remove the risk, by stopping particular contractors working at their premises.

#### Enforcement

The requirements of HSWO are enforceable through improvement and prohibition notices. Improvement notices give a time limit for compliance with requirements. Prohibition notices can have the effect of immediately stopping the operations of a business or the activities of a person where imminent risk is apparent. Failure to comply with HSWO may result in court action.

#### AGE LIMITS AND CONSENT

The need for limits on age, and requirements for consent, can be a controversial area because in many circumstances specific requirements have not been made in law. Consent is a complex area of law, and one that is often misunderstood by the general public, and also by some skin piercing practitioners.

#### Tattooing

The [Tattooing of Minors \(Northern Ireland\) Order 1979](#) imposes a statutory minimum age of 18 years for permanent tattooing (except when carried out for medical reasons by a duly qualified medical practitioner or by a person working under their direction). The practitioner has a defence if they can show that they had good reason to believe that the person was over 18 years of age. The consent of a client under 18 is not a defence. The Police enforce this legislation and fines are up to £500.

#### Skin piercing activities

There is no statutory age of consent for cosmetic piercing (cosmetic body piercing and ear piercing). Cosmetic piercing of a minor is lawful provided a valid consent is given. Furthermore the courts have held that a parents right to decide on behalf of his and her child yields to the child's competence to make a decision, for example if he or she is capable of understanding the nature of the act to be done (see below). Body piercing for sexual gratification is unlawful. Children under the age of 17 are not able to consent lawfully to a piercing that would be regarded as indecent assault. Genital or nipple piercing performed on someone under the age of 17 might be regarded as an indecent assault under sexual offences legislation depending on the facts of the case.

Ear piercing and in some cultures nose piercing is generally considered acceptable when carried out on a minor, even below the age of five, provided that a parent or legal guardian gives consent and is present whilst the procedure is carried out

Click on text to view

**SUMMARY****Specific controls****General controls****Determining requirements****REGISTRATION AND BYELAW REQUIREMENTS IN NORTHERN IRELAND****HEALTH AND SAFETY AT WORK (NI) ORDER 1978****General duties****Risk assessment****Control of substances****hazardous to health****Management of contractors****Enforcement****AGE LIMITS AND CONSENT****Tattooing****Other skin piercing activities**

Acupuncture and electrolysis

**Consent****USE OF LOCAL ANAESTHETIC MEDICATION**

# Tattooing and body piercing guidance

## PART A

### Legislative background on tattooing and skin piercing activities (*continued*)

#### Acupuncture and electrolysis

When carried out properly these do not cause harm to the body, and leave no permanent markings. For this reason they are not likely to raise any concerns over common assault charges, although practitioners should be aware of possible indecent assault complications (see below). Consent should still be obtained before treatment takes place, and in the case of a minor this should be obtained from the parents or legal guardians.

#### Other considerations

Semi-permanent skin-colouring, cosmetic piercing, beading, branding, scarring, cutting and other extreme forms of body modification do cause actual harm and generally leave permanent marks and can result in disfigurement. They can therefore be considered as assaults to the body, and so potentially subject to the legislation concerning assault. This means that the question of age and the client's informed consent are very important

#### Consent

In the legal proceedings of *R v Brown* (1994) 1 AC 212, the House of Lords ruled on appeal that consent could not be a defence against sections 20 and 47 of the Offences Against the Person Act 1861 which deals with common assaults. However, the law also recognises that certain activities that give rise to 'harm' are lawful. This includes surgery, tattooing, ear piercing and violent sports. The courts have also held that the law allows children under the age of 18 to consent to cosmetic body piercing provided they are sufficiently mature to understand the nature of the request. This kind of assessment is clearly a subjective matter for the operator who will need to ensure that the client is provided with sufficient information to allow them to proceed in an informed way and without pressure.

Under the Sexual Offences (Northern Ireland) Order 2008, girls and boys under the age of 16 cannot legally give consent to intimate sexual contact under any circumstances, so piercing of nipples and genitalia (for girls) or genitalia (for boys) can be regarded as an assault offence. Evidence that such contact was for sexual gratification would be required in order to constitute an indecent assault. The Female

Genital Mutilation Act 2003 states that certain procedures in respect of female genitals are illegal unless carried out for medical reasons.

The signing of a declaration and providing proof of age should be a fundamental part of the client consultation process and practitioners should always require that the client signs a consent form prior to any work being commenced. However, the consent will only be valid if the customer has been fully informed as to the nature of the process, the likely effect and potential problems involved. An example of a consent form is provided in Appendix 8. However, for practitioners own protection, it is recommended that any consent forms they use are worded with the advice of a solicitor who is familiar with this area of law.

#### USE OF LOCAL ANAESTHETIC MEDICATION

There is a range of topical local anaesthetic products for surface (skin) anaesthesia available from community pharmacies (Pharmacy only (P) medicines), however none of these products are licensed for local anaesthesia prior to tattoo or body piercing. The client may wish to obtain a topical local anaesthetic preparation prior to the procedure; however responsibility for purchasing and application of the product should remain with the client. The client should be advised to read the Patient Information Leaflet which accompanies the product and should be aware of the following:

- Warnings, cautions and contraindications.
- Side effects.
- That they are using the licensed product for an un-licensed indication.
- Recommendations regarding administration and application.

Alternatively, a qualified practitioner, e.g. doctor, may prescribe a topical local anaesthetic product to be self-administered by the client, or can prescribe and administer a topical local anaesthetic product, in accordance with legal requirements of their professional registration.

Local anaesthetic injections are prescription-only medicines (POMs) therefore they can only be prescribed by a suitably qualified practitioner. Local anaesthetic injections are not licensed for



## Section 01

Click on text to view

**SUMMARY****Specific controls****General controls****Determining requirements****REGISTRATION AND  
BYELAW REQUIREMENTS IN  
NORTHERN IRELAND****HEALTH AND SAFETY AT  
WORK (NI) ORDER 1978****General duties****Risk assessment****Control of substances****hazardous to health****Management of contractors****Enforcement****AGE LIMITS AND CONSENT****Tattooing****Other skin piercing activities**

Acupuncture and electrolysis

**Consent****USE OF LOCAL ANAESTHETIC  
MEDICATION**

# Tattooing and body piercing guidance

## PART A

### Legislative background on tattooing and skin piercing activities (*continued*)

local anaesthesia prior to tattoo or body piercing. In addition they should not be administered parenterally unless adequate resuscitation equipment is available. Information on the supply and administration of injectable medicines outside their licensed medicinal uses is available from the Medicines and Healthcare Regulatory Agency (MHRA)<sup>2</sup> and from the Nursing and Midwifery Council (NMC).

**References**

1. British National Formulary (BNF) <http://www.medicinescomplete.com/mc/bnf/current/PHP8663-local-anaesthesia.htm> (accessed 22/4/2013)
2. MHRA; Frequently asked questions: Supply and administration of Botox®, Vistabel®, Dysport® and other injectable medicines outside their licensed medicinal uses such as in cosmetic procedures <http://www.mhra.gov.uk/Howweregulate/Medicines/Availabilityprescribingandsupplyingofmedicines/Frequentlyraisedissues/BotoxVistabelDysportandotherinjectablemedicinesincosmeticprocedures/index.htm> (accessed 22/4/2013)
3. NMC; Remote prescribing and injectable cosmetic medicinal products <http://www.nmc-uk.org/Nurses-and-midwives/Regulation-in-practice/Regulation-in-Practice-Topics/Remote-prescribing-and-injectable-cosmetic-medicinal-products/> (accessed 22/4/2013)
4. DHSSPSNI; Local Government (Northern Ireland) Order 2005 Regulation of Cosmetic Piercing and Skin-Colouring businesses Guidance on Article 31 and Schedule 2 <http://www.dhsspsni.gov.uk/bodypiercing-order.pdf> (accessed on 20/08/14)



# Tattooing and body piercing guidance

## PART A

### Section 02a Infection prevention and control

Click on text to view

**SECTION 02A**

**INTRODUCTION**

**Infection, its causes and spread**

**Factors in infection control practice**

**Responsibilities**

**STANDARD PRINCIPLES OF INFECTION CONTROL**

**Appendix 01 – Infection, its causes and spread**

**SECTION 02B**

**PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE**

**SECTION 02C**

**PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT**

**SECTION 02D**

**PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS**

**SECTION 02E**

**PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS**

**SECTION 02F**

**PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT**

**SECTION 02G – REFERENCES**

### INTRODUCTION

#### Infection, its causes and spread

The causes and methods of spread of infections that are likely to arise in connection with tattooing and skin piercing, are well understood.

#### See Appendix 01– Infection, its causes and spread

Unsafe or unhygienic practices by tattooing/body piercing practitioners can lead to the spread of infectious diseases that can affect the health of the client as well as jeopardise the health of the practitioner. Although some bacterial or viral infections may be spread during procedures that do not involve skin penetration, it is the occupational risk of transmission of infections such as blood-borne viruses (BBVs), such as hepatitis B, hepatitis C, hepatitis D and HIV, which can arise and which are of primary concern. Precautions to minimise the possibility of exposure to blood from an infected client or practitioner should be put in place by the adoption of safe practices and procedures. This should include immunisation against hepatitis B (Department of Health, 2010a).

#### Factors in infection control practice

The risk of transmission of infection can be minimised by:

- Good cleanliness of the premises where the tattooing and/or body piercing is taking place, and of the fixtures and fittings.
- Good personal hygiene of the practitioners.
- Correct cleaning and sterilization or disposal of instruments, materials and equipment processes in place.

It is therefore important that the safe working practices described in this guidance are followed at all times in order to protect both the client and practitioner.

### Responsibilities

Under the HSWO, all employers should ensure that all their employees are appropriately trained and proficient in the procedures necessary for working safely. Employers and their employees are also responsible to ensure that any person on the premises is not placed at any avoidable risk, as far as is reasonably practicable. They also have a responsibility to protect voluntary workers. Employers are also required by COSHH, to review every procedure carried out by their employees which involves contact with a substance hazardous to health, including pathogenic micro-organisms. Specific guidance is available from the Department of Health (Department of Health, 1998).

### STANDARD PRINCIPLES OF INFECTION CONTROL

This guidance is based upon standard principles which are the basic level of infection control practice. Compliance with these standard principles reduces the risk of transmission of blood-borne and other pathogens.

Everyone providing treatments to clients should know about and be able to carry out these standard principles for infection prevention and control (National Institute for Health and Care Excellence 2012). To that end they should have received training in:

- Hand hygiene and skin care.
- The use of personal protective equipment (PPE).
- Sharps management and management of exposure to blood and body fluids.
- Safe handling, storage and disposal of waste materials.
- Cleaning and disinfection of the environment.

# Tattooing and body piercing guidance

## PART A

### Section 02b Infection prevention and control

Click on text to view

**SECTION 02A**

**INTRODUCTION**

**STANDARD PRINCIPLES OF INFECTION CONTROL**

**SECTION 02B**

**PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE**

**Hand washing facilities**

**When to wash hands**

**What to use to wash hands**

**How to carry out hand washing**

**Use of hand rubs**

**Hand care**

Use of hand cream

Care of broken skin

**Poster – How to handwash**

**SECTION 02C**

**PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT**

**SECTION 02D**

**PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS**

**SECTION 02E**

**PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS**

**SECTION 02F**

**PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT**

**SECTION 02G – REFERENCES**

### PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE

Hand hygiene is a major component of the standard principles and one of the most effective methods to prevent transmission of pathogens by reducing the number of microorganisms that may be present. The spread of infection from hands is well recognised and the importance of compliance with hand hygiene practices is emphasised in all national and international guidelines.

#### Hand washing facilities

Hand washing facilities should be adequate and conveniently located in treatment areas. Hand washing instructions should be clearly displayed at the hand wash basin, such as in the form of a poster.

#### See Poster – How to handwash

Hand wash basins must be designated for that purpose only and have a constant supply of hot and cold running water, ideally delivered through a mixer tap. Under no circumstances should equipment be washed in hand wash basins.

Liquid soap dispensers with single use liquid soap cartridges/bottles should be used, ideally wall-mounted, although free standing dispensers would be considered suitable. Disposable liquid soap cartridges are recommended because they do not permit a topping-up process and this minimizes the risk of contamination.

Wall-mounted disposable paper towels should be next to the hand wash basins, and fully stocked at the start of each working day to minimise or reduce the need to fill up within hours during which the premises is operational.

A foot-operated pedal bin, of an appropriate size, should be placed next to the hand wash basin for disposal of paper towels.

As a gold standard for infection control purposes, it is recommended that hand wash basins should:

- Have elbow/foot-operated or non-touch mixer taps.
- Have wall-mounted cartridge soap dispensers and paper towels available at each hand wash basin.
- Not have a plug or overflow or be capable of taking a sink plug.
- Not have taps aligned to run directly into the drain aperture.
- Have waterproof splashbacks.
- Have space allowed at the design stage for the placement of waste bins next to the hand wash basin.

#### When to wash hands

- Before and after an intervention with each client.
- After contact with any blood or body fluids.
- Immediately after the removal of gloves.
- After using a tissue or handkerchief.
- After smoking.
- After visiting the toilet.
- Before and after eating.
- Immediately after any other activity or contact with a client's surroundings that could potentially result in hands becoming contaminated.

#### What to use for hand washing

For an ordinary hygienic hand wash, the use of liquid soap is sufficient. Preparations containing antiseptics that have a residual effect on the skin surface are not required for use in tattoo and body piercing settings.

## Section 02b Infection prevention and control

Click on text to view

## SECTION 02A

## INTRODUCTION

## STANDARD PRINCIPLES OF INFECTION CONTROL

## SECTION 02B

## PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE

## Hand washing facilities

## When to wash hands

## What to use to wash hands

## How to carry out hand washing

## Use of hand rubs

## Hand care

Use of hand cream

Care of broken skin

## Poster – How to handwash

## SECTION 02C

## PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT

## SECTION 02D

## PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS

## SECTION 02E

## PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS

## SECTION 02F

## PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT

## SECTION 02G – REFERENCES

**How to carry out hand washing**

There are three distinct and essential stages to handwashing (National Institute for Health and Care Excellence, 2012)

**1. Preparation**

Before washing hands, all wrist and hand jewellery should be removed. Cuts and abrasions must be covered with waterproof dressings. Fingernails should be kept short, clean and free from nail polish. Hands should be made wet by placing them under tepid running water before applying liquid soap.

**2. Washing and rinsing**

The hand wash solution must come into contact with all of the surfaces of the hand. The hands must be rubbed together vigorously for a minimum of 10–15 seconds, paying particular attention to the tips of the fingers, the thumbs and the areas between the fingers. Hands should be rinsed thoroughly.

**3. Drying**

In a tattooing and body piercing setting, good quality disposable soft paper towels would be considered the method of choice because communal towels are a source of cross-contamination. Paper towels should be stored in a wall-mounted dispenser next to the washbasin and thrown away in a pedal operated waste bin. Hands should not be used to lift the lid or they will become re-contaminated.

**Use of hand rubs**

Hand rubs containing alcohol based products can enable practitioners to quickly and effectively clean their hands before and after contact with clients. However, the use of a hand alcohol rub/gel is not a substitute for using soap and water for hand washing e.g. when undertaking tattooing and body piercing procedures (see above), and should not be used when the hands are visibly soiled or potentially contaminated with body fluids (National Institute for Health and Care Excellence, 2012). Hand rubs should conform to the standard BS EN 1500. The hand rub solution must come into contact with all surfaces of the hand; the hands must be rubbed together paying particular attention to the tips of the fingers, the thumbs and the areas between the fingers, until the solution has evaporated.

**Hand care****Use of hand cream**

A hand cream can be applied regularly to protect skin from the drying effects of regular hand decontamination (National Institute for Health and Care Excellence, 2012). Each practitioner should have their own supply and a communal pot should not be used.

**Care of broken skin**

Unbroken skin is the best defence because it provides the perfect barrier against infection. Small areas of broken or infected skin on exposed parts of the practitioner's body should be covered with a waterproof dressing that completely covers the affected area.

# Tattooing and body piercing guidance

## PART A

### Section 02c Infection prevention and control

Click on text to view

**SECTION 02A**

**INTRODUCTION**

**STANDARD PRINCIPLES OF INFECTION CONTROL**

**SECTION 02B**

**PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE**

**SECTION 02C**

**PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT**

**Assessment of risk**

**Types of protective clothing**

Work clothing

Gloves

- Glove choice
- Synthetic materials
- Latex gloves
- Sterile gloves
- Gloves used for cleaning

Aprons

Eye and face protection

**SECTION 02D**

**PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS**

**SECTION 02E**

**PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS**

**SECTION 02F**

**PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT**

**SECTION 02G – REFERENCES**

↪ Main contents

### PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT

Personal protective equipment (PPE) should be available to all practitioners and staff who may be at risk whilst working in the premises.

#### Assessment of risk

Regulation 3 of the Management of Health and Safety at Work Regulations (Northern Ireland) 2000 requires every employer to make a suitable and sufficient assessment of:

- a) risks to the health and safety of their employees to which they are exposed whilst they are at work; and
- b) risks to health and safety of persons not in their employment arising out of or in connection with the conduct by them of their undertaking.

Therefore the selection of protective equipment must be based on an assessment of the risk of transmission of infection between the practitioner and client and vice versa:

Anticipated level of exposure	Wear disposable gloves	Wear plastic or fluid repellent apron	Wear eye and face protection
No exposure to blood/body fluids anticipated	X	X	X
Exposure to blood/body fluids anticipated but low risk of splashing	Yes	Yes	X
Exposure to blood/body fluids anticipated with high risk of splashing to the face	Yes	Yes	Yes

### Types of protective clothing

#### Work clothing

Practitioner clothing should be clean at all times, and professional in appearance. Work clothing should be changed daily. Staff clothing should not impede good hand washing, therefore the wearing of short sleeved tops is advocated (Department of Health, 2010b).

#### Gloves

COSHH requires employers to assess any substances hazardous to health, including biohazards within blood and body fluids (such as blood-borne viruses) and take steps to reduce the risk of exposure.

The use of gloves has two purposes:

1. To protect the hands from becoming contaminated with dirt and microorganisms.
2. By changing gloves, to prevent transfer of microbes from one client to another.

Gloves must be worn when carrying out invasive procedures, when in contact with sterile sites and non-intact skin or mucous membranes, and during all activities that have been assessed as carrying a risk of exposure to blood, body fluids, secretions or excretions, or to sharp or contaminated instruments (National Institute for Health and Care Excellence, 2012).

The correct method for wearing and removing gloves can be found on the WHO Save Lives website: ([http://www.who.int/gpsc/5may/Glove\\_Use\\_Information\\_Leaflet.pdf](http://www.who.int/gpsc/5may/Glove_Use_Information_Leaflet.pdf)). Hands must be washed and dried thoroughly before putting on disposable gloves.

Gloves can tear or puncture visibly during use, or leakage may occur through microscopic holes. Hands may also become contaminated as gloves are removed. Gloves therefore must not be seen as a substitute for good hand hygiene. Used gloves should be disposed of as offensive waste (see waste section).

## Section 02c Infection prevention and control

Click on text to view

**SECTION 02A****INTRODUCTION****STANDARD PRINCIPLES OF INFECTION CONTROL****SECTION 02B****PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE****SECTION 02C****PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT****Assessment of risk****Types of protective clothing**

## Work clothing

## Gloves

- Glove choice
- Synthetic materials
- Latex gloves
- Sterile gloves
- Gloves used for cleaning

## Aprons

## Eye and face protection

**SECTION 02D****PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS****SECTION 02E****PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS****SECTION 02F****PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT****SECTION 02G – REFERENCES**

Gloves are single-use items (National Institute for Health and Care Excellence, 2012). This means they must be put on immediately before an episode of client contact or procedure and removed as soon as the activity is completed, or when leaving the client for any reason. Gloves must be changed between different procedural activities for the same client and between dealing with different clients, or removed if they become torn.

**Glove choice**

Gloves should be made available in a range of sizes for use by different practitioners (National Institute of Clinical Excellence, 2012).

All gloves used for direct client care must conform to current EU legislation (CE marked as for single use) and should be appropriate for the task. Only PPE meeting the basic health and safety requirements of the EC Personal Protective Equipment Directive requirements is entitled to carry a CE mark and be sold for use in the EC (Health and Safety Executive, 2012a). Practitioners should therefore look for the CE mark information on glove packs plus EN 374-1:2003 or EN 374-2. These markings show the gloves are protective against chemicals and can resist microorganisms at a performance level 2 test in penetration tests. Although this cannot infer protection against viruses, because they are not used in the performance tests, in practice this is the highest level of protection afforded against microorganisms (Health and Safety Executive, 2012a).

**Synthetic materials**

Neoprene and nitrile gloves are synthetic gloves which have been shown to have comparable in-use barrier performance to natural rubber latex gloves in laboratory and clinical studies.

**Nitrile**

- Provides an excellent biological barrier, resistant to punctures and tears.
- Comparable to latex in terms of barrier performance.
- Is a good alternative for latex sensitive individuals.
- Can be used where a latex free environment is necessary.
- Is less elastic than latex but does shape to the wearer's hand over time.
- Can be used for handling certain chemicals (Infection Control Nurses Association, 2002).

However, nitrile contains the same types of chemicals as latex in the manufacturing process and allergic reactions have been reported.

**Polyisoprene and Neoprene**

- Offers effective protection against viral penetration.
- Has similar elasticity and physical properties as latex.
- Is suitable for individuals sensitised to latex proteins.
- Can be used when a latex free environment is necessary (Infection Control Nurses Association, 2002).

**Vinyl**

- Is suitable for use in areas where there is a low biohazard risk.
- Provides a good alternative for use when staff or clients are sensitised to latex.
- In lab tests shows increased permeability to blood borne viruses than latex.
- Possesses lower tensile strength than latex and breaks down more frequently.
- Is prone to leaking.
- Is inelastic and can be baggy to wear.
- Is inexpensive in comparison to synthetic rubbers (Infection Control Nurses Association, 2002).



## Section 02c Infection prevention and control

Click on text to view

**SECTION 02A****INTRODUCTION****STANDARD PRINCIPLES OF INFECTION CONTROL****SECTION 02B****PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE****SECTION 02C****PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT****Assessment of risk****Types of protective clothing**

## Work clothing

## Gloves

- Glove choice
- Synthetic materials
- Latex gloves
- Sterile gloves
- Gloves used for cleaning

## Aprons

## Eye and face protection

**SECTION 02D****PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS****SECTION 02E****PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS****SECTION 02F****PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT****SECTION 02G – REFERENCES**

Overall, vinyl gloves can be used to perform many tasks, but, depending on the quality of the glove may not be appropriate when handling blood/blood-stained fluids (Royal College of Nursing, 2012).

**Polythene**

These gloves are not recommended for use when undertaking activities involving blood/body fluids exposure and therefore should not be used in the tattooing and body piercing setting. They are ill fitting, have heat sealed seams that are predisposed to split and have a tendency to tear (Infection Control Nurses Association, 2002).

**Latex gloves (natural rubber latex)**

Latex gloves may be the preferred choice for procedures in tattooing and body piercing because of latex's tactile sensitivity, barrier property against viruses, good fit and optimal elasticity and user familiarity. However, latex is a known skin and respiratory sensitiser and in a small number of people it can cause serious allergy (see below) (Health and Safety Executive, 2012a). There may also be issues when using latex gloves alongside petroleum-based lubricants which may affect the glove's integrity and therefore its protection ability.

Latex allergies are becoming common with prolonged use of latex gloves (Infection Control Nurses Association, 2002). The use of appropriate synthetic gloves is therefore recommended to avoid becoming sensitised. It is recognised, however, that within certain work environments, latex gloves are still used in large numbers due to their efficacy and low cost. If latex gloves are worn, then powder free, low protein content materials must be chosen to help prevent latex allergy (Health and Safety Executive, 2012).

Where latex gloves are in use, monitoring of clients and staff should be undertaken. Any sensitivity shown to natural rubber latex in either clients or staff should be documented and action must be taken to remove further exposure risks. Alternatives to natural rubber latex gloves therefore must be made available (Health and Safety Executive, 2012). Further information on latex allergy can be found on-line at: <http://www.hse.gov.uk/skin/employ/latex-gloves.htm> and detailed information on skin care and dermatitis in the work place can be found at: <http://www.hse.gov.uk/skin/>

**Sterile gloves**

Sterile gloves are used for major surgical procedures and are not applicable to tattooing/body piercing.

**Gloves used for cleaning**

For environmental cleaning purposes or for manual pre-cleaning of equipment prior to disinfection/sterilization, general-purpose rubber gloves should be used. The gloves should be washed with general-purpose detergent and warm water, and dried between uses. They should be changed weekly, or more frequently if the gloves become damaged (for example if there are signs of peeling, cracking and tears) (National Patient Safety Agency, 2007).



# Tattooing and body piercing guidance

## PART A

### Section 02d Infection prevention and control

Click on text to view

#### SECTION 02A

##### INTRODUCTION

##### STANDARD PRINCIPLES OF INFECTION CONTROL

#### SECTION 02B

##### PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE

#### SECTION 02C

##### PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT

#### SECTION 02D

##### PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS

###### Introduction

Sharps and needles

Sterile needles

###### 'Sharps/needlestick' injuries and exposure to blood/body fluid spills

Types of injury/exposure

Risks of transmission of blood-borne viruses following a sharps injury/exposure

Management of sharps/needlestick injuries and exposure to blood and body fluids

###### Sharps/'needlestick' injuries

###### Blood/body fluid spills

###### Occupational health for blood borne virus prevention

Risk assessment

Vaccination requirements

###### Appendix 02 – Blood borne viruses

###### Appendix 03– Safe use and disposal of sharps

###### Appendix 04– First Aid following a sharps/ 'needlestick' injury and exposure to blood/body fluid

###### Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill

#### SECTION 02E

##### PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS

#### SECTION 02F

##### PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT

#### Section 02g – References

↪ Main contents

#### Aprons

A disposable plastic apron must be worn when there is a risk that clothing may be exposed to blood, body fluids, secretions or excretions (with the exception of sweat or tears) (National Institute for Health and Care Excellence, 2012).

Plastic aprons should be used as single-use items and changed between clients. They should be discarded and disposed of as offensive waste after use.

#### Eye and face protection

Eye protection and face masks must be worn where there is a risk of blood, body fluids, secretions or excretions splashing into the eyes and face (National Institute for Health and Care Excellence, 2012). A risk assessment of the planned procedure should be undertaken to help inform decision making (e.g. when manually cleaning equipment as part of decontamination processes).

If reusable goggles/protective glasses are used, they should be washed after each client or task using a general purpose detergent, rinsed and stored dry. Eye protection should be compatible with any facemask used.

Face masks (such as surgical masks) should only be used if there is a risk of splashing of blood/body fluid droplets into the mouth or nose. If used, masks should be changed between clients and disposed of immediately after use. They must not be carried or worn around the neck.

## PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS

### Introduction

All body fluids should be regarded as potentially infectious. Blood carries the highest risk of transmitting blood borne viruses such as hepatitis B, C, D and human immunodeficiency virus (HIV). Blood borne viruses may also be transmitted by other body fluids, especially if contaminated by blood (Health Protection Agency, 2009).

### See appendix 02 – Blood borne viruses

#### Sharps and needles

The word 'sharps' is a generic term that includes needles, scalpels, stitch cutters, glass ampoules and sharp instruments that may become contaminated with blood or body fluid. In tattooing/body piercing premises, sharps include equipment such as razors, needle bars with needles attached and cannulae (sometimes used for body piercing).

'Sharps' contaminated with blood or other body fluids should be classified as hazardous waste and handled accordingly.

### See Section 02e

All 'sharps' must be handled and disposed of safely and with extreme care. After use they should be placed immediately into yellow sharps boxes/bins with orange lids, compliant with UN 3291 and BS7320 standards. This is to reduce the risk of exposure to blood-borne viruses, for example through an accidental 'sharps' or 'needlestick' injury (National Institute for Health and Care Excellence, 2012).

### See Appendix 03 – Safe use and disposal of sharps

#### Sterile needles

Only sterile single-use needles should be used for skin piercing or tattooing. Needles should be examined for imperfections prior to their use and discarded if any are found. Needles should either be used directly from the packaging or placed on a sterile surface/tray for immediate use.

# Tattooing and body piercing guidance

## PART A

### Section 02d Infection prevention and control

Click on text to view

#### SECTION 02A

#### INTRODUCTION

#### STANDARD PRINCIPLES OF INFECTION CONTROL

#### SECTION 02B

#### PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE

#### SECTION 02C

#### PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT

#### SECTION 02D

#### PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS

##### Introduction

Sharps and needles

Sterile needles

##### Sharps/needlestick injuries and exposure to blood/body fluid spills

Types of injury/exposure

Risks of transmission of blood-borne viruses following a sharps injury/exposure

Management of sharps/needlestick injuries and exposure to blood and body fluids

##### 'Sharps/needlestick' injuries

##### Blood/body fluid spills

##### Occupational health for blood borne virus prevention

Risk assessment

Vaccination requirements

#### Appendix 02 – Blood borne viruses

#### Appendix 03– Safe use and disposal of sharps

#### Appendix 04– First Aid following a sharps/needlestick injury and exposure to blood/body fluid

#### Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill

#### SECTION 02E

#### PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS

#### SECTION 02F

#### PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT

#### Section 02g – References

↪ Main contents

### 'Sharps/needlestick' injuries and exposure to blood and body fluids

#### Types of injury/exposure

A blood/body fluid injury/exposure incident includes:

- Inoculation of blood by a needle or other 'sharps'.
- Contamination of broken skin with blood.
- Blood splashes to mucous membrane, e.g. eyes or mouth.
- Swallowing a person's blood, e.g. after mouth-to-mouth resuscitation.
- Contamination where the individual has an open wound, and clothes have been soaked by blood.
- Bites (where the skin is broken).

#### Risks of transmission of blood-borne viruses following a significant injury/exposure

Transmission of blood borne viruses (BBVs) may result from contamination of mucous membranes of the eyes or the mouth, or of broken skin, with infected blood or other infectious material. There is no evidence that BBVs can be transmitted by blood contamination of intact skin, inhalation or by faecal-oral contamination.

The transmission risks after a **mucocutaneous** exposure (splash exposure) are lower than those after a **percutaneous** exposure ('sharps' injury), estimated at 1 in a 1000 for HIV (Health Protection Agency 2008). There is currently no evidence on the risk of transmission for hepatitis B virus (HBV) and hepatitis C virus (HCV) following mucocutaneous exposure (Health Protection Agency, 2008).

The risk of infection following a percutaneous injury, especially deep penetrating injuries involving a hollow-bore needle or a device visibly contaminated with blood has been estimated at:

- 1 in 3 when a source patient is infected with HBV and is classed as being highly infectious at the time.
- 1 in 30 when the patient is infected with HCV.
- 1 in 300 when the patient is infected with HIV (Health Protection Agency, 2008).

#### Management of sharps/needlestick injuries and exposure to blood and body fluids

Injuries where a person's broken skin or eyes, mouth or other mucous membranes are exposed to another person's blood or body fluids, therefore, may carry a risk of infection with blood borne viruses.

#### Sharps/needlestick injuries

Prompt first aid and immediate risk assessment is needed in the event of such incidents to establish the type of exposure sustained and to help determine what appropriate action is needed.

#### See Appendix 04 – First Aid following a blood/body fluid exposure

#### Blood/body fluid spills

Blood and body fluid spills must be dealt with quickly and effectively. Specialist body fluid spill kits are available to purchase. These can be stored in a safe designated area in the premises, enabling easy access and timely clear up. The expiry dates of products inside kits should be regularly checked and out of date items replaced as necessary. Posters and simple training should be provided on the use of the body fluid spill kits.

# Tattooing and body piercing guidance

## PART A

### Section 02d Infection prevention and control

Click on text to view

#### SECTION 02A

##### INTRODUCTION

##### STANDARD PRINCIPLES OF INFECTION CONTROL

#### SECTION 02B

##### PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE

#### SECTION 02C

##### PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT

#### SECTION 02D

##### PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS

##### Introduction

Sharps and needles

Sterile needles

##### Sharps/ 'needlestick' injuries and exposure to blood/ body fluid spills

Types of injury/exposure

Risks of transmission of blood-borne viruses following a sharps injury/exposure

Management of sharps/ needlestick injuries and exposure to blood and body fluids

##### Sharps/ 'needlestick' injuries

##### Blood/body fluid spills

##### Occupational health for blood borne virus prevention

Risk assessment

Vaccination requirements

##### Appendix 02 – Blood borne viruses

##### Appendix 03– Safe use and disposal of sharps

##### Appendix 04 – First Aid following a sharps/ 'needlestick' injury and exposure to blood/body fluid

##### Appendix 05 – Protocol for cleaning up a blood or a blood stained body fluid spill

#### SECTION 02E

##### PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS

#### SECTION 02F

##### PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT

#### Section 02g – References

↪ Main contents

The body fluid spill kit should contain:

- Disposable plastic aprons and synthetic (e.g. nitrile) gloves.
- Disposable cloths.
- General purpose detergent.
- Chlorine granules.

**See Appendix 05 – Protocol for cleaning up blood or a blood stained body fluid spill**

### Occupational health for blood borne virus prevention

#### Risk assessment

Apart from the overall duty to carry out risk assessment of hazards in the workplace, COSHH places a specific duty on employers to assess the risks from exposure to hazardous substances, including pathogens (called biological agents in COSHH), and to bring into effect the measures necessary to protect workers and others from those risks as far as is reasonably practicable.

In these circumstances the assessment of risks to health should include:

- How to prevent exposure to biological agents.
- Steps needed to achieve adequate control of exposure.
- Steps needed to avoid accidental 'needlestick/sharps' injury.

#### Vaccination requirements

Those at risk of blood/body fluid exposure through sharps or splashes, therefore, should have a full course of hepatitis B vaccine. An accelerated course consisting of three doses at zero, one and two months (followed by a fourth dose at twelve months after the first dose for those at continued risk of exposure), and antibody titres (blood levels) should be checked one to four months after the completion of the primary course of vaccine. It is recommended that those at continued risk of infection should be offered a once only single booster, approximately five years after completion of the primary immunisation course (antibody levels do not need to be checked before or after this booster dose). (Department of Health 2006).

Under the HSWO, employers must pay for protective measures such as immunisation. This is usually provided through the company occupational health provider. In the absence of an occupational health service, the employee could be asked to arrange immunisation through their own GP, but the employer must make alternative arrangements if this cannot be done, and reimburse any charges made to the employee for such arrangements. As with all control measures, immunisation needs to be checked and reviewed and boosters provided, where necessary.

It would be considered good practice for practitioners to keep copies of their vaccination history/antibody level results. If practitioners refuse to have hepatitis B vaccination, it is advised the employer should consider asking the employee to sign a disclaimer form.

If the response to the hepatitis vaccine is not sufficient, the GP will need to investigate whether there is a specific reason for non-response to the vaccine. It is most important for non-responders to know their status. They may need to be protected by other measures (e.g. Immunoglobulin) following a 'needlestick/sharps' injury.

There is no vaccine against hepatitis C and human immunodeficiency virus (HIV). Robust infection control measures should be employed at all times to minimise the risk of exposure to these viruses.

If a practitioner is found to be positive for a blood borne virus disease, they should be assessed and advised by their GP in relation to working practices.

# Tattooing and body piercing guidance

## PART A

### Section 02e Infection prevention and control

Click on text to view

**SECTION 02A**

**INTRODUCTION**

**STANDARD PRINCIPLES OF INFECTION CONTROL**

**SECTION 02B**

**PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE**

**SECTION 02C**

**PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT**

**SECTION 02D**

**PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS**

**SECTION 02E**

**PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS**

**Responsibility for waste materials**

**National guidance on waste management**

Offensive waste

Sharps

All other waste

Disposal of aerosol cans, glass, bottles, broken crockery and dry cell batteries

**Appendix 06 – Principles for good waste handling**

**SECTION 02F**

**PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT**

**Section 02g – References**

↪ Main contents

### PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS

Good waste management is important to:

- Reduce the health and safety risk to staff, clients and visitors.
- Protect the environment.
- Reduce waste disposal costs.

#### Responsibility for waste materials

All organisations have a legal responsibility to dispose of waste safely, ensuring no harm is caused either to staff, members of the public or the environment. This responsibility begins when waste is generated and ends with its final disposal. It is essential that persons handling waste fulfil their legal responsibilities by taking care to prevent injury or transmission of infection to themselves or others.

Premises should have a waste policy. The practitioner is responsible for ensuring that contracts are in place for collection and safe disposal of offensive/hazardous waste from the premises. It is essential to ensure with the waste management provider that appropriate documentation is generated when necessary. The manager of the premises is also responsible for monitoring the performance of staff and waste contractors, as per agreed contract.

#### National guidance on waste management

In 2013 The Department of Health, Social Services and Public Safety published Health Technical Memorandum HTM 07-01- The “Safe Management of Healthcare Waste” ([http://www.dhsspsni.gov.uk/htm\\_07-01\\_final.pdf](http://www.dhsspsni.gov.uk/htm_07-01_final.pdf)). This document is to be used as a best practice guide for the safe and effective handling of waste. The previous clinical waste classification system using groups A to E can no longer be used, as the groups do not reflect the appropriate segregation for transport or disposal.

#### See Appendix 06– Principles for good waste handling

There are a few municipal non-healthcare waste streams that are classed to be similar in nature to healthcare waste and are hence included within the Department of Health’s waste guidance’s assessment framework. Waste such as ‘sharps’ and related wastes from tattoo and body piercing practice is specifically included in this definition (Department of Health 2012).

Soft waste from body-piercing or tattooing practice can normally be assumed to present no risk of infection, unless an indication to the contrary is provided by a healthcare professional (Department of Health 2012). However, as waste contaminated with non-infectious body fluids is capable of causing offence, it would be classed as ‘offensive’ waste (see below) and would require appropriate packaging to indicate the bag contents (Department of Health 2012). Sharps waste would always be considered hazardous waste and should be disposed of accordingly (see below) (Department of Health 2012).

#### Offensive waste

The guidance from the Department of Health, Social Services and Public Safety states that “used gloves and aprons, swabs, small dressings, and cotton wool contaminated with body fluids arising from cosmetic piercing and other body art plus other special treatment procedures would be considered as offensive/ hygiene waste where it is generated in quantity (one bag of 7 kg or more in any collection interval)”. Offensive waste should be placed into a yellow/black bag (“tiger bag”) for disposal so that subsequent holders of the waste can handle and dispose of the material appropriately. Only where such waste “is generated in small quantities (less than 7 kg in one bag in any collection period) should it be disposed in the black-bag (general waste) stream” (Department of Health 2012).

## Section 02e Infection prevention and control

Click on text to view

**SECTION 02A****INTRODUCTION****STANDARD PRINCIPLES OF INFECTION CONTROL****SECTION 02B****PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE****SECTION 02C****PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT****SECTION 02D****PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS****SECTION 02E****PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS****Responsibility for waste materials****National guidance on waste management**

Offensive waste

Sharps

All other waste

Disposal of aerosol cans, glass, bottles, broken crockery and dry cell batteries

**Appendix 06 – Principles for good waste handling****SECTION 02F****PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT****Section 02g – References****Sharps'**

'Sharps' should always be handled and disposed of as hazardous waste ([see Section 02d](#)).

In Northern Ireland 'sharps' not contaminated with medicinal products (such as those generated at a body piercing/tattooing premises), should be disposed of in a standards compliant yellow 'sharps' bin with an orange lid (Department of Health, 2012).

**All other waste**

All other non-contaminated waste such as paper should be placed in black bags, or bags for recycling, within a foot operated pedal bin and disposed of as normal household waste. Aerosols, batteries and broken glass should not be placed in these bags.

**Disposal of Aerosol Cans, Glass, Bottles, Broken Crockery and Dry Cell Batteries**

These items should always be placed in a designated cardboard box, lined with a plastic bag so that it is leak-proof. The box should be labelled to indicate its contents and method of disposal.



## Section 02f Infection prevention and control

Click on text to view

**SECTION 02A****INTRODUCTION****STANDARD PRINCIPLES OF INFECTION CONTROL****SECTION 02B****PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE****SECTION 02C****PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT****SECTION 02D****PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS****SECTION 02E****PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS****SECTION 02F****PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT****Equipment for cleaning****Colour-coding for cleaning equipment****Use of chemicals****Appendix 07 – Template protocol for environmental cleaning of premises****Section 02g – References****PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT**

Cleaning is the process that physically removes contamination with organic material such as blood and body fluids, along with dirt and dust. Cleaning does not necessarily destroy microorganisms from the item that is being cleaned. However, providing and maintaining a clean and readily cleanable environment facilitates the prevention and control of infections.

**Equipment for cleaning**

Cleaning equipment that is regularly used should be fit for purpose, easy-to-use and well-maintained (National Patient Safety Agency, 2007a). A clutter-free environment and the adoption of local 'clean as you go' policies will provide the foundation for quality service provision in a clean, safe place. Good cleaning practice includes having:

- The provision and maintenance of a clean and appropriate environment by using systems to manage the environment's cleanliness with a documented cleaning policy and rota plus a regular audit programme.
- Equipment which is suitable for purpose, is able to be kept clean (i.e. impervious surfaces) and maintained in good physical repair.
- Working from the cleanest area towards the dirtiest area to greatly reduce the risk of cross contamination.
- A person in charge who has direct responsibility for ensuring that cleanliness standards are maintained.
- Single use cloths for cleaning tasks and cleaning equipment such as mops and buckets kept in good order (i.e. cleaned daily, renewed regularly and stored safely (cleaned, dried and stored inverted) in a designated area after use).

**See Appendix 07 – Template protocol for environmental cleaning of premises**

**Colour-coding for cleaning equipment**

The aim of a colour-coding system for cleaning equipment is to prevent cross-contamination. There is a national colour-coding system for the NHS (National Patient Safety Agency, 2007b) (e.g. blue for general areas, red for toilet areas/wash hand basins), which could be adapted for use in the tattooing and body piercing setting.

**Use of chemicals**

Household detergent is adequate for most routine environmental cleaning. For high risk environmental surfaces such as treatment surfaces, a hypochlorite solution of 1000 parts per million (PPM) available chlorine should be used. This solution should be made up for use on a daily basis, following the manufacturer's instructions, in a labelled container provided by the commercial manufacturer (using, for example, one tablet of sodium dichloroisocyanurate (NaDCC) per litre format). After twenty-four hours the solution must be discarded. The hypochlorite solution must not be transferred into a trigger spray bottle but be used directly from the container onto a disposable cloth or paper towels. Surfaces contaminated with blood should be cleaned in accordance with the guidance on dealing with blood spillage as a higher concentration of hypochlorite will be required (**see Appendix 05 – Protocol for cleaning up blood or a blood stained body fluid spill**).

All chemicals should be handled and stored in accordance with the manufacturer's instructions/COSHH guidance (Health and Safety Executive, 2002). Material safety data sheets should be accessible to all staff. All chemicals used on the premises should be used and stored in an identified cool, dry and well ventilated place (room/cabinet) that is lockable, out of reach of visitors and members of the public and in the original containers. Expiry dates should be routinely checked.



## Section 02g Infection prevention and control

Click on text to view

**SECTION 02A****INTRODUCTION****STANDARD PRINCIPLES OF INFECTION CONTROL****SECTION 02B****PRINCIPLES OF INFECTION CONTROL – HAND HYGIENE****SECTION 02C****PRINCIPLES OF INFECTION CONTROL – PERSONAL PROTECTIVE EQUIPMENT****SECTION 02D****PRINCIPLES OF INFECTION CONTROL – MANAGEMENT OF SHARPS AND EXPOSURE TO BLOOD AND BODY FLUIDS****SECTION 02E****PRINCIPLES OF INFECTION CONTROL – SAFE HANDLING, STORAGE AND DISPOSAL OF WASTE MATERIALS****SECTION 02F****PRINCIPLES OF INFECTION CONTROL – CLEANING AND DISINFECTION OF THE ENVIRONMENT****Section 02g – References****References**

Calderdale Local Authority (2008). Advice and safe practice for permanent tattooing plus advice and safe practice for body piercing. (<http://www.calderdale.gov.uk/business/licences/skinpiercing/index.html>)

Dartford Borough Council (2009). Code of practice for hygienic piercing. ([www.dartford.gov.uk](http://www.dartford.gov.uk))

Department of Health (1998). UK Health Departments Guidance for Clinical Health Care Workers: Protection Against Infection with Blood-borne Viruses Recommendations of the Expert Advisory Group on AIDS and the Advisory Group on Hepatitis. HMSO ([http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_4002766](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4002766))

The Department of Health, Social Services and Public Safety, Health Technical Memorandum HTM 07-01- The Safe Management ([http://www.dhsspsni.gov.uk/htm\\_07-01\\_final.pdf](http://www.dhsspsni.gov.uk/htm_07-01_final.pdf))

Department of Health (2006). Immunisation against Infectious Disease. ([http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_079917](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_079917))

Department of Health (2010b). Uniforms and workwear: guidance on uniform and workwear policies for NHS employers. ([http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_114751](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_114751))

Department of Health (2012). Safe management of healthcare waste V2.0. ([http://www.dh.gov.uk/prod\\_consum\\_dh/groups/dh\\_digitalassets/documents/digitalasset/dh\\_133874.pdf](http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/digitalasset/dh_133874.pdf))

Department of Health, Social Services and Public Safety, Health Technical Memorandum HTM 07-01- The Safe Management of Health Care Waste. ([http://www.dhsspsni.gov.uk/htm\\_07-01\\_final.pdf](http://www.dhsspsni.gov.uk/htm_07-01_final.pdf))

Health Protection Agency (2008). Eye of the Needle: UK surveillance of significant exposures to BBV in healthcare workers. ([www.hpa.org.uk/web/HPAwebFile/HPAweb\\_C/1227688128096](http://www.hpa.org.uk/web/HPAwebFile/HPAweb_C/1227688128096))

Health Protection Agency (2009). Inoculation Injuries and Children in Schools and similar settings: Risk Assessment Guidelines for Health Protection Units. ([http://www.hpa.org.uk/webc/HPAwebFile/HPAweb\\_C/1259152291335](http://www.hpa.org.uk/webc/HPAwebFile/HPAweb_C/1259152291335))

Health and Safety Executive (2002). Guidance on Control of Substances Hazardous to Health Regulations. (<http://www.hse.gov.uk/coshh/>)

Health and Safety Executive (2008). Health and Safety Regulation – a short guide. (<http://www.hse.gov.uk/pubns/hsc13.pdf>)

Health and Safety Executive (2012a). The use of gloves. Blood Borne Viruses Guidance. (<http://www.hse.gov.uk/biosafety/blood-borne-viruses/use-of-gloves.htm>)

Health and Safety Executive (2012b). How to deal with an exposure incident. Blood Borne Viruses Guidance. (<http://www.hse.gov.uk/biosafety/blood-borne-viruses/how-deal-exposure-incident.htm>)

Health & Safety at Work etc Act (1974). (<http://www.legislation.gov.uk/ukpga/1974/37>)

Infection Control Nurses Association (ICNA) (now Infection Prevention Society (IPS)) (2002). A comprehensive glove choice. Fitwise, England.

Management of Health and Safety at Work Regulations. ([www.opsi.gov.uk/si/si1999/19993242.htm](http://www.opsi.gov.uk/si/si1999/19993242.htm))

National Institute for Health and Care Excellence (2012). Infection control: Prevention and control of healthcare-associated infections in primary and community care (CG139). (<http://guidance.nice.org.uk/CG139>)

National Patient Safety Agency (NPSA) (2007a). The national specifications for cleanliness in the NHS: a framework for setting and measuring performance outcomes. (<http://www.nrls.npsa.nhs.uk/resources/?EntryId45=59818>)

National Patient Safety Agency (NPSA) (2007b). NHS colour coding hospital cleaning materials and equipment 15. (<http://www.nrls.npsa.nhs.uk/resources/?EntryId45=59810>)

National Patient Safety Agency (NPSA) (2009). The Revised Healthcare Cleaning Manual. (<http://www.nrls.npsa.nhs.uk/resources/?EntryId45=61830>)

Royal College of Nursing (2012). Wipe it out- Essential practice for infection prevention and control. Guidance for nursing staff. ([http://www.rcn.org.uk/\\_data/assets/pdf\\_file/0008/427832/004166.pdf](http://www.rcn.org.uk/_data/assets/pdf_file/0008/427832/004166.pdf))

## Tattooing and body piercing guidance

### PART A

#### Section 03 Before and aftercare of a tattoo or body piercing

Click on text to view

**SKIN PREPARATION**

**NEED FOR AFTERCARE**

**Care of skin after tattooing**

**Care of skin after piercing**

**Aftercare leaflets**

**WHAT TO DO IF A CLIENT RETURNS WITH AN INFECTION**

**Appendix 08– Tattooing and body piercing consent form**

**Appendix 09– Aftercare follow-up record sheet**

**References**

#### SKIN PREPARATION

Where shaving is required, only single-use razors are acceptable. Either a 70% alcohol-impregnated single use swab (typically 70% isopropyl alcohol) or a 0.5% chlorhexidine in 70% alcohol single use swabs (National Institute for Health and Care Excellence, 2012) can be used for skin disinfection. If the skin is visibly dirty then the area should first be cleaned with soap and water and dried with a paper towel.

Where it is necessary to mark the skin, a single use toothpick dipped into gentian violet or other suitable dye could be used. The dye should be dispensed into a single-use pot for each client. Otherwise the entire bottle should be discarded after each client. As an alternative, or where large areas of skin need to be marked, then a single-use commonly available marker pen could be used, or a suitable single-use alternative. (Dartford Borough Council, 2009)

Where products such as petroleum jelly are used for procedures, an appropriate amount of material should be dispensed, using a single-use implement (Calderdale Local Authority, 2008), into a single-use pot for every client. Practitioners should not use cream/lotion direct from a jar/tube. Roll-on or stick applicators are not acceptable for use. The practitioner's hands, even if gloved, should never come into contact with the contents of these jars/tubes.

#### NEED FOR AFTERCARE

All piercings must be performed using good infection prevention technique and pre-sterilized equipment to reduce the chance of infection occurring. The aftercare of body piercing sites and tattooed areas is also very important in order to promote good healing which lowers the risk of infection and reduces the risk of scarring.

Practitioners must explain to their clients the known potential complications associated with the particular procedure they are being asked to carry out. Upon completion of a treatment, they must provide both verbal and written aftercare advice relevant to that treatment.

In the case of body piercing of minors, they must have a parent or other responsible adult present when aftercare advice is given.

Good practice is to include on the consent form a tick box or similar indicator to record that aftercare advice has been explained and discussed at the time that the consent form was signed.

**See Appendix 08 – Tattooing/body piercing consent form**

Maintaining a good level of hygiene around the treated area is essential during the healing period.

#### Care of skin after tattooing

Good practice is to cover the tattooed area with sterile non-adhesive gauze which is then secured with hypo-allergenic tape. Gauze permits ventilation and aids healing.

A sterile, non-adhesive dressing may be appropriate for larger areas, at least during the client's journey home, but in many cases simply keeping the area clean and dry is likely to be the best approach. If plastic film wrap is used for larger areas then it must be clean (taken directly from the pack) and the client should be advised on when and how to replace this covering.

#### Care of skin after piercing

The piercing site should be kept clean and dry to promote healing.

When checking the pierced site, hands should be clean. It is not normally necessary to rotate or remove a piece of jewellery however, if the client has been advised to do this, it should be handled as little as possible, with clean hands and using a clean tissue and, if possible, to touch only the jewellery item.

Clients should be advised about healing times as these may be prolonged because of the time it takes for the jewellery "tunnel" to become dry and then to heal after the initial piercing.

## Tattooing and body piercing guidance

### PART A

#### Section 03 Before and aftercare of a tattoo or body piercing

Click on text to view

**SKIN PREPARATION**

**NEED FOR AFTERCARE**

**Care of skin after tattooing**

**Care of skin after piercing**

**Aftercare leaflets**

**WHAT TO DO IF A CLIENT RETURNS WITH AN INFECTION**

**Appendix 08 – Tattooing and body piercing consent form**

**Appendix 09– Aftercare follow-up record sheet**

**References**

#### Aftercare leaflets

Lapses in aftercare are common causes of infection following piercing and clients should be provided with an appropriate aftercare leaflet such as those available within this guidance.

Click on the aftercare leaflet thumbnail below to view the links for downloading

#### 01 Tattoo



#### 02 Ear and face piercing



#### 03 Oral piercing



#### 04 Body and surface piercing



#### 05 Genital piercing (female)



#### 06 Genital piercing (male)



#### 07 Microdermal implants



#### WHAT TO DO IF A CLIENT RETURNS WITH AN INFECTION

If any of the following signs or symptoms of an infection develop, urgent medical attention should be sought:

- Redness spreading around the site and extending away from it.
- Pus or green/yellow fluid oozing from the site.
- Bleeding that is not controlled by light pressure.
- Pain (rather than discomfort).
- Swelling.
- Heat.
- Immobility of, or reluctance to move, a limb/digit/part of the body.

Where the client informs the practitioner of a concern or problem, the practitioner is advised to keep records of any action taken and advice given.

See Appendix 09– Aftercare follow-up record sheet

#### References

Calderdale Local Authority (2008). Advice and safe practice for permanent tattooing plus advice and safe practice for body piercing. (<http://www.calderdale.gov.uk/business/licences/skinpiercing/index.html>)

Dartford Borough Council (2009). Code of practice for hygienic piercing. ([www.dartford.gov.uk](http://www.dartford.gov.uk))

National Institute for Health and Care Excellence (2012). Infection control: Prevention and control of healthcare-associated infections in primary and community care (CG139). (<http://guidance.nice.org.uk/CG139>)

# Tattooing and body piercing guidance

## PART A

### Section 04 Decontamination

Click on text to view

**PRINCIPLES OF DECONTAMINATION**

[Need for vigilance](#)

[Categorization of risks](#)

**DECONTAMINATION PRACTICE**

[Layout of decontamination area](#)

[Decontamination of invasive items](#)

[Cleaning](#)

[Use of disinfectants](#)

[Sterilization](#)

[Sterilizer operation, validation, maintenance and record keeping](#)

[Care of items after sterilization](#)

[Decontamination of blood and body fluids in the working environment items](#)

[Appendix 10– Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard: self-assessment and decision making tool for tattoo and body piercing practitioners](#)


[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13 – Autoclave daily record sheet](#)

[Further reading](#)

### PRINCIPLES OF DECONTAMINATION

Decontamination is a combination of processes that render reusable items safe for reuse.

- For invasive items used in tattooing and body piercing, these decontamination processes will involve cleaning and sterilization, after which environmental recontamination should be minimised and recontamination with blood must be totally eliminated. Pre-sterilized single-use items are a good alternative.
- Using items once and then discarding them removes the need for decontaminating them.
- Where an item is marked  it is single-use and must not be reused even if decontaminated.
- For items used in association with invasive items, these decontamination processes should include both cleaning and sterilization. Where this is not possible (tattooing motors for example), their contamination should be minimised by the use of impervious covers which should be removed carefully after use so as not to contaminate the surface of the item, and the item itself should be chemically disinfected. Alternatively items can be single-use, such as with ink caps. As with invasive items, environmental recontamination should be minimised and recontamination with blood must be totally eliminated.
- For surfaces, contamination with blood should be made safe by careful application of chemical disinfectants.

#### Need for vigilance

It is vital that, once decontaminated, items are not directly or indirectly contaminated with blood or body fluid. This requires scrupulous handling procedures and physical separation from undecontaminated items and the surfaces they may have contaminated.

### Categorization of risks

A general categorization of the risks that items pose with regard to transmission of infection and the minimum decontamination standard that should apply is set out in the following table:

Risk	Use of item	Decontamination (minimum standard)
High	Items introduced into normally sterile body areas or in contact with a break in skin or mucous membrane	Sterile or sterile single-use
Medium	In contact with intact mucous membranes	Disinfect or single-use
Low	In contact with intact skin	Clean or clean and disinfect if contaminated with blood or body fluid
Minimal	Not normally in contact with skin (e.g. floors and walls)	No specific treatment required, domestic cleaning. Spills or splashes of blood or body fluid should be safely cleared.

### DECONTAMINATION PRACTICE

#### Layout of decontamination area

The layout of a decontamination facility is important, whether it is in a separate room or a dedicated part of the treatment room, within body art premises. Items to be decontaminated must flow along a defined process pathway from dirty (i.e. used and contaminated), through cleaning (which may have both a manual and an ultrasonic stage), through sterilization and into a phase of clean storage and return to use.



Click on text to view

**PRINCIPLES OF  
DECONTAMINATION**

**Need for vigilance**

**Categorization of risks**

**DECONTAMINATION PRACTICE**

**Layout of decontamination  
area**

**Decontamination of invasive  
items**

Cleaning

Use of disinfectants

Sterilization

Sterilizer operation, validation,  
maintenance and record keeping

Care of items after sterilization

**Decontamination of blood and  
body fluids in the working  
environment items**

**Appendix 10 –  
Decontamination requirements  
for equipment used in  
tattooing and skin piercing**

**Appendix 11– Equipment  
sterilization standard: self-  
assessment and decision making  
tool for tattoo and body piercing  
practitioners**

**Appendix 12 – Equipment  
and body piercing jewellery  
sterilization standard for  
tattooists and body piercers**

**Appendix 13 – Autoclave daily  
record sheet**

**Further reading**

↪ Main contents

At any stage, items must not be re-contaminated by direct or indirect contact with items at a lesser stage of decontamination, e.g. items that have been cleaned must not be put on the same surface or handled with the same utensils used for dirty items; items that have been sterilized must not make contact with any surface that has been used for items before they have been sterilized. In addition to this beware of using a clean-looking item in the mistaken belief that it has been fully decontaminated.

Only by having a progression through a defined layout, can the location of an item correspond to its stage in the decontamination process. This is always important, but particularly so if more than one practitioner uses the same facility, or if multiple copies of identical equipment items are available for use, e.g. tattoo machine grips.

It is an advantage if the decontamination occurs in a dedicated room but if this is not possible, it should still occur in a dedicated area in which a defined flow from dirty to clean can be clearly established.

### Decontamination of invasive items

Any item that pierces the skin poses a high risk of transmission of infection. Only items intended for reuse after cleaning and steam sterilization should be reused. If any uncertainty exists whether an item can be decontaminated by this method then confirmation should be obtained from the supplier.

#### See Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing

#### Cleaning

All items should be cleaned before sterilization. If any blood or other proteinaceous material is left on an item that is to be steam sterilized it will become firmly fixed on the items and very difficult to remove subsequently.

Cleaning should use methods, detergents and concentrations of those detergents compatible with items and specifically intended for instrument cleaning. The cleaning method should not put practitioners at risk from contaminants on

instruments: cleaning should be done under the detergent solution surface to prevent splashing and care should be taken to avoid injury.

Any detergent should be safe for those who use it, but thick washing-up gloves should nevertheless be used. If manual cleaning is used, the detergent should be at or around neutral pH. It is common to use detergents containing enzymes. However, it is thought that short exposures of dirt containing proteins to these enzymes may give insufficient time for effective action. There are also concerns that some of the enzyme mixtures (“subtilisins”) can give rise to allergic reactions in some who come into contact with them. This should be considered in risk assessments. Use of non-enzymatic detergents should be considered.

Cleaning should be carried out in a sink dedicated to instrument cleaning and not one shared with other functions (handwashing, eating utensil washing etc.). It can help to attain the correct dilution of a detergent if the sink is indelibly marked to a fill-line of known volume, allowing a measured addition of detergent to achieve the correct dilution. Warm, but not hot, water should be used (hot water may coagulate proteins onto an item and make then difficult to remove). Utensils used for instrument cleaning should be dedicated solely for that purpose and should be observably fit for purpose (e.g. bristles on brushes in good order). Cleaning should take place under the surface of the detergent solution to minimise the potential of splashing the practitioner with the blood being removed.

Cleaning should be done as soon after an item is used as possible as drying makes contamination less easily removed. It may make cleaning easier if items are stored between use in clean fresh tap water or in a high humidity atmosphere (such as a closed vessel with a small amount of water).

Validation of cleaning should be by careful visual inspection of each item in good light.

Items with complex surfaces may require subsequent additional cleaning in an ultrasonic waterbath (ultrasound creates



## Section 04 Decontamination

Click on text to view

**PRINCIPLES OF  
DECONTAMINATION**

**Need for vigilance**

**Categorization of risks**

**DECONTAMINATION PRACTICE**

**Layout of decontamination  
area**

**Decontamination of invasive  
items**

Cleaning

Use of disinfectants

Sterilization

Sterilizer operation, validation,  
maintenance and record keeping

Care of items after sterilization

**Decontamination of blood and  
body fluids in the working  
environment items**

**Appendix 10 –  
Decontamination requirements  
for equipment used in  
tattooing and skin piercing**

**Appendix 11 – Equipment  
sterilization standard: self-  
assessment and decision making  
tool for tattoo and body piercing  
practitioners**

**Appendix 12 – Equipment  
and body piercing jewellery  
sterilization standard for  
tattooists and body piercers**

**Appendix 13– Autoclave daily  
record sheet**

**Further reading**

strong microcurrents in water by a process known as “cavitation” and is very effective at removing soiling from otherwise hidden crevices). Ultrasonic waterbaths should be used, maintained and validated according to the manufacturer’s instructions.

**Use of disinfectants**

Chemical disinfection has far lower levels of quality assurance than steam sterilization and should not be used for invasive items. It should only be used for decontamination of the environment and non-invasive items. Chemical disinfectants are usually inactivated by organic matter and should only be used after cleaning has removed the vast majority of organic matter. Disinfectants should be used in a controlled manner according to guidance and manufacturer’s instructions (e.g. the correct dilution, freshly prepared, applied as directed). Do not mix disinfectants with other detergents or chemicals unless following manufacturer’s instructions.

The disinfectant of choice for general disinfection of the environment should be a hypochlorite solution containing 1,000 parts per million available chlorine (ppm av Cl). This is usually made by dilution of tablets of sodium dichloroisocyanurate (NaDCC), a form of solid, stable hypochlorite, pre-measured to give specific hypochlorite concentration when dissolved in given volumes of water (always follow manufacturer’s instructions for attaining the correct concentration). Whilst solid NaDCC is stable on dry storage, the hypochlorite solutions it generates are unstable and should be made-up daily. Always use in accordance with the manufacturer’s materials safety data sheet. Hypochlorite solutions may bleach fabrics and corrode metals other than good quality stainless steel. They should never be mixed with strong acids (the production of highly toxic chlorine gas can result). They should be used in well ventilated areas.

**Sterilization**

Sterilization is the complete elimination of all microbial life to a very high level of quality assurance. It must be a robustly efficient process guaranteed to work on every occasion. Whilst there are many theoretical ways in which sterilization can be achieved, the only way that body art practitioners can achieve it with the required quality is by steam sterilization.

Other methods are either too complex or lack equivalent quality assurance. Steam sterilizers are devices that can expose items to be sterilized to pure steam at above atmospheric pressure in a chamber. The process must be automatic and steam sterilizers must monitor the process to ensure that all parameters of sterilization have been met or, if any have failed, to clearly indicate a failed cycle. (Pressure cookers do not have this inbuilt quality assurance and are not suitable for body art instrument sterilization. Devices sold as baby bottle steam sterilizers use lower temperatures and are not suitable for body art instrument sterilization). The type of steam sterilizer suitable for use by body art practitioners is known variously as a small steam sterilizer, a benchtop steam sterilizer or a transportable steam sterilizer. These are small steam sterilizers that generate their own steam and are powered by a standard domestic electrical supply and should conform to the standard BS EN 13060.

To effect the energy transfer necessary for sterilization, steam must be able to condense on all surfaces of an item, therefore hinged items should be opened and items should not overlap each other. If pockets of air are present, this blocks the ability of steam to make contact with the surface in air. If porous (e.g. fabrics), hollow or wrapped items are placed in chambers that are then filled with steam, this will not remove air from these loads but will only compress the air in them, leading to inadequate sterilization.

## Section 04 Decontamination

Click on text to view

**PRINCIPLES OF DECONTAMINATION**

Need for vigilance

Categorization of risks

**DECONTAMINATION PRACTICE**

Layout of decontamination area

Decontamination of invasive items

Cleaning

Use of disinfectants

Sterilization

Sterilizer operation, validation, maintenance and record keeping

Care of items after sterilization

Decontamination of blood and body fluids in the working environment items

Appendix 10 –

Decontamination requirements for equipment used in tattooing and skin piercing

Appendix 11– Equipment sterilization standard: self-assessment and decision making tool for tattoo and body piercing practitioners

Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers

Appendix 13 – Autoclave daily record sheet

Further reading

- Small steam sterilizers are produced in 3 different types:

Type B steam sterilizers (also known as “vacuum” or “porous load” sterilizers): These have sterilization cycles that start by pulling a vacuum on the chamber to remove most of the air in the chamber and within porous, hollow or wrapped items; this is followed by a series of pulses and removal of steam to dilute any remaining air, before the chamber is brought to sterilizing conditions. These sterilizers are suitable for porous, hollow or wrapped items. Any wrapping must conform to the standard BS EN 868, showing they are of a type that does not prevent the passage of steam and will resist the passage of contamination after sterilization has occurred.

Type S steam sterilizers: These are similar to type B, but have only been validated to process specific loads (and are normally produced to sterilize wrapped dental handpieces). These sterilizers are suitable for porous, hollow or wrapped items only if validated for those specific items to be processed.

Type N steam sterilizers: These are steam sterilizers with no assisted air removal. These sterilizers are suitable for non-porous, non-hollow (solid) and unwrapped items only.

Type B and S sterilizers tend to be more expensive to buy and maintain and normally have longer cycle times than type N.

**See Appendix 11– Equipment sterilization standard: self-assessment and decision making tool for tattoo and body piercing practitioners**

**See Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers**

### **Sterilizer operation, validation, maintenance and record keeping**

Sterilizers should only be used by those trained in their correct operation. Sterilizers should be operated according to the sterilizer manufacturer’s instructions.

A steam sterilizer should monitor each sterilization cycle and produce a record (e.g. printout) of the cycle parameters, primarily the temperatures attained throughout the sterilizing phase and the times of those temperature readings. If the sterilizer cannot produce a record of cycle parameters, equivalent records should be produced by observation of a cycle at the start of each day the sterilizer will be used. Steam sterilizers should be validated and maintained by people specifically trained to do so according to a schedule provided by the sterilizer manufacturer. Records of validation and maintenance should be retained as locally advised by the inspecting authority.

**See Appendix 13 – Autoclave daily record sheet**

There is a legal requirement to have a written scheme of periodic examination of steam sterilizers under the Pressure Systems Safety Regulations (Northern Ireland) 2004. The written scheme shall be drawn up by, or certified as being suitable by a competent person as defined by the Regulations. A certificate of insurance is required for the sterilizer as a pressure vessel (Medical Devices Agency, 2002).

### **Care of items after sterilization**

If items are wrapped, they will remain sterile as long as the wrapping remains intact and dry.

If items are unwrapped, they can be placed in a clean, lidded container. Great care should be taken not to recontaminate them, with particular emphasis on recontamination from undecontaminated instruments or surfaces contaminated by blood or body fluid as they are transferred into or out of the container, or as other items are removed from the container. This is best achieved by having only those items in one container that will be used in a single procedure. If a container does become contaminated, it should be washed and processed (open) in a steam sterilizer or discarded.

## Section 04 Decontamination

Click on text to view

**PRINCIPLES OF DECONTAMINATION**

Need for vigilance

Categorization of risks

**DECONTAMINATION PRACTICE**

Layout of decontamination area

Decontamination of invasive items

Cleaning

Use of disinfectants

Sterilization

Sterilizer operation, validation, maintenance and record keeping

Care of items after sterilization

**Decontamination of blood and body fluids in the working environment items****Appendix 10– Decontamination requirements for equipment used in tattooing and skin piercing****Appendix 11 – Equipment sterilization standard: self-assessment and decision making tool for tattoo and body piercing practitioners****Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers****Appendix 13 – Autoclave daily record sheet****Further reading****Decontamination of blood and body fluids in the working environment**

Blood and other body fluids in the environment pose a risk of infection transmission: with blood the main hazards are of blood borne viruses; with other body fluids a wider spectrum of infectious micro-organisms are relevant. Environmental contamination with blood or other body fluids should be dealt with by the removal of the contaminating material and disinfection as soon as possible after the contamination occurs. The longer any contamination remains on a surface, the more it poses a direct risk to those who come in contact with it, as well as the indirect risk of it being transferred to other surfaces from which it may contaminate instruments that come into contact with clients.

Remember: The person most at risk, is the person clearing the contamination. They should have safe methods of working and use appropriate personal protective equipment (PPE) – which should always include single-use gloves, with additional PPE such as aprons if the contamination is extensive.

Most environmental contamination is likely to be minor. Such contamination can be removed using single-use wipes and then, once clean, the area treated with a suitable disinfectant such as hypochlorite. (See cautions on hypochlorites above). If there is likely to be more extensive contamination, purpose-made disinfectant spill kits can be brought-in in readiness and used.

**Further reading**

Medicines and Healthcare products Regulatory Authority (MHRA). Sterilization, Disinfection and Cleaning of Medical Equipment: Guidance on Decontamination from the Microbiology Advisory Committee to Department of Health. Part 1 – Principles (2010); part 2 – Protocols (2005); Part 3 – Procedures (2006). <http://www.mhra.gov.uk/Publications/Safetyguidance/Otherdevicesafetyguidance/CON007438>

Medical Devices Agency (2002). Benchtop Steam Sterilizers – Guidance on Purchase, Operation and Maintenance.

(<http://www.mhra.gov.uk/groups/dts-bi/documents/publication/con007327.pdf>)

Hoffman PN, Bradley CR, Ayliffe GAJ (2004). Disinfection in Healthcare. 3rd Edn. Blackwell Publishing. ISBN 1405126426

## Section 05 Product quality of tattoo ink

Click on text to view

**BACKGROUND AND CONTEXT****MICROBIOLOGICAL AND CHEMICAL QUALITY CONCERNS****RECOMMENDATIONS TO IMPROVE THE QUALITY OF INK PRODUCTS USED FOR TATTOOING****References****BACKGROUND AND CONTEXT**

Requirements already exist at both European and national level, whereby ink products should be sterile and inert at first use. These specified standards are not necessarily legally binding, though bylaw requirements would become so if adopted by a local authority. At the European level the quality requirements for inks were initially laid down in 2003 by the Council of Europe, and further revised in 2008 (see at – <https://wcd.coe.int/ViewDoc.jsp?id=45869&Lang=en>).

A common requirement for byelaws is that the inks should be sterile at first use and should be inert (i.e. non-reactive with the body). The containers used to hold the inks for each customer should be either disposed of at the end of each session of treatment, or cleaned and sterilized before re-use.

For EHOs out in the field and for tattooists, there is an expectation that ink manufacturers will provide products that are fit for purpose. However, at present there are few independent sources of data to confirm the quality of inks used in the UK, and no authoritative information to indicate whether this is universally achievable. Some studies have shown that certain products may be contaminated with microorganisms and/or metals, and the quality of inks used in the UK is likely to vary between manufacturers because of an absence of common standard quality requirements.

**MICROBIOLOGICAL AND CHEMICAL QUALITY CONCERNS**

The risks associated with tattooing treatments have been acknowledged for many years, and have been reported internationally (Lehman et al., 2010). In addition to the standard safer sharps controls to mitigate the risk of Blood Borne Virus (BBV) transmission during tattooing, another potential hazard is that of infection via the 'environmental' route. This is typically associated with naturally occurring bacteria and fungi that have gained entry to the ink product at some stage during its manufacture or storage.

This form of environmental contamination of inks, prior to their use on the client, is rarely reported on. Despite the continually improving standards in tattooing health and safety - much of it related to eliminating the risk of BBV transmission - the 'environmental' aspect of tattooing infection control remains beyond the control of most practitioners.

Poor quality tattoo inks increase the potential for localised bacterial skin infections as well as dermal allergies following tattooing treatments (Limentani et al., 1979; Jacob, 2002; Charnock, 2004; Suhair, 2007; Drage et al., 2010). Ink related problems might therefore be directly related to the chemical and/or microbiological quality of the ink or pigments used. Where this occurs, other efforts to maintain tattooing hygiene standards could potentially be undermined if the inks themselves are contaminated at the point of use.

Tattoo ink products are typically purchased from suppliers or directly from manufacturers and are delivered intra-dermally during treatment, so there is an increased potential for the client's body to be exposed to their components, compared with, for example, a topically applied skin colorant.

Some inks are sold with little or no accompanying product data, and their composition may remain uncertain even at the point of use.

Reports of metal sensitivity following tattooing have been published in the UK, and dermatological responses have been linked with particular ink colours, especially red products [some containing mercury] and greens/blue [chromate/cobalt] (Jacob, 2002; Mortimer et al., 2003; Gass & Todd, 2007).

**RECOMMENDATIONS TO IMPROVE THE QUALITY OF INK PRODUCTS USED FOR TATTOOING:**

- The presence of an accompanying product data sheet is a fundamental requirement to ensure that, as far as possible, the appropriate quality tests have been undertaken and passed for the ink. Tattooists should therefore work wherever

## Section 05 Product quality of tattoo ink

Click on text to view

**BACKGROUND AND CONTEXT**

**MICROBIOLOGICAL AND CHEMICAL QUALITY CONCERNS**

**RECOMMENDATIONS TO IMPROVE THE QUALITY OF INK PRODUCTS USED FOR TATTOOING**

**References**

possible with inks that are accompanied by the manufacturer's product quality information, and should request this information from their supplier if it is not provided; and,

- Practitioners should note the batch numbers of the products they purchase, with delivery dates, as new inks are purchased and received. Some suppliers already provide a product listing and batch numbers with their delivery note, and this would equally serve as a dated record of ink products received. This record can then be used in case of any subsequent concerns over ink quality. Wherever possible, and to strengthen the quality control link between client and product(s) used, the colour of the inks used on each client should be noted at the time of treatment.

**References**

Charnock C (2004). Tattooing dyes and pigments contaminated with bacteria. *Tidsskr Nor Laegeforen.* 124(7):933-5.

Drage LA, Ecker PM, Orenstein R, Phillips PK and Edson RS (2010). An outbreak of *Mycobacterium chelonae* infections in tattoos. *Journal of the American Academy of Dermatology.* 62(3):501-6.

Gass JK and Todd PM (2007). Multiple manifestations of chromate contact allergy. *Contact Dermatitis.* 56:290–291.

Jacob CI (2002). Tattoo-associated dermatoses: a case report and review of the literature. *Dermatol Surg.* 28(10):962-5.

Lehman EJ, Huy J, Levy E, Viet SM, Mobley A and McCleery TZ (2010).

Bloodborne pathogen risk reduction activities in the body piercing and tattooing industry. *American Journal of Infection.* 38(2):130-8.

Limentani AE, Elliott LM, Noah ND, Lamborn JK (1979). An outbreak of hepatitis B from tattooing. *Lancet.* 2(8133):86-8.

Mortimer NJ, Chave TA and Johnston GA (2003). Red tattoo reactions. *Clinical and Experimental Dermatology.* 28:508–510.

Suhair OS, Miles HFJ, Simmons P, Stickley J and De Giovanni JV (2007). Awareness of the risk of endocarditis associated with tattooing and body piercing among patients with congenital heart disease and paediatric cardiologists in the United Kingdom. *Archives of Disease in Childhood.* 92:1013-1014



## Section 06 Body piercing jewellery

Click on text to view

[GENERAL](#)[NICKEL DIRECTIVE](#)[References](#)**GENERAL**

The style of body piercing jewellery is distinct from traditional jewellery worn in the ear lobe such as studs and butterfly designs.

Variations on the barbell and (captive) ring design are those most commonly seen in body piercing because these substantial shapes minimise the risk of embedding, tearing and migration.

High quality jewellery is made with smooth surfaces and joins in order to reduce the risk of irritation or of harbouring infection.

Practitioners should ensure that all jewellery used for skin piercings is sterile prior to its use. Where jewellery is not purchased pre-sterilized but is sterilized within the premises, the method by which sterilization has been carried out will determine how sterile the jewellery remains (see sterilization section). If jewellery is processed and stored properly, it may retain its sterility indefinitely. However, practitioners should be advised to discuss issues such as shelf life with equipment (and packaging) manufacturers.

A number of materials are used in skin piercing jewellery and acceptable materials include:

- Titanium.
- Niobium.
- Platinum.
- Gold – preferably solid gold 14 carat or 18 carat (for ear piercing). The use of gold higher than 18 carat is not recommended in body piercing as it is too soft and the potential exists for scratching or pitting of the metal which may increase the risk of infection at the piercing site. The use of gold lower than 14 carat is not recommended in body piercing as it tends to be lower in quality and has the potential to contain metallic impurities, which may lead to allergic response in the pierced individual.

**NICKEL DIRECTIVE**

The Nickel Directive was a European Union Directive regulating the use of nickel in jewellery and other products that come into contact with the skin. The requirements also cover a wide range of other items such as necklaces, bracelets, wristwatch cases, zips and buttons.

Since June 2009 it has been subsumed into the EU REACH Regulation (Registration, Evaluation, Authorisation, legislation and restriction of Chemicals). Nevertheless, the term nickel directive is still used to refer to the restrictions on nickel usage and the prescribed test for quantifying nickel release from products. Jewellery can only be used if the nickel release rate from those parts of these products coming into direct and prolonged contact with the skin is 0.5 micrograms per square centimetre per week, or less. For body piercing jewellery, post assemblies - the part of the jewellery that is inserted into the wound caused by the piercing of the skin, including both the piece that goes through the wound and those parts of the jewellery intended to hold the piece in and against the wound (earring “back” or balls on the end of a piercing bar or stud) - are prohibited unless their rate of nickel release is 0.2 micrograms per square centimetre per week, or less.

One problem with these requirements is that they do not apply to jewellery manufactured for export to countries outside the European Union. Where practitioners cannot prove that jewellery being used is in compliance with these requirements, it is advised that use of that jewellery stops until the practitioner can obtain evidence from the manufacturer of its compliance.

**References**

The Regulation now incorporates the Nickel Directive (REACH), at item 27 of Annex XVII.

Further information is available from: <http://www.teg.co.uk/nickel/94-27-EC.htm>

# Tattooing and body piercing guidance

## PART A

### Section 07 Governance

Click on text to view

**TRAINING AND COMPETENCIES**

**PROCEDURE MANUALS/ POLICIES**

**AUDIT AND QUALITY MONITORING**

**RECORD KEEPING**

**CONSENT AND AFTERCARE DOCUMENTATION**

See Appendix 08– contains an example of a consent form and there are a variety of written aftercare leaflets available

See Appendix 09– Aftercare follow-up record sheet

**References**

#### **TRAINING AND COMPETENCIES**

All tattooists and body piercers should undertake accredited training and be able to demonstrate their competencies before starting to practice. This guidance recognises, however, that tattooists and body piercers in the UK do not have access to a nationally recognised and accredited course, nor is there a nationally agreed set of competencies or ongoing objective monitoring of professional competence by a recognised professional body.

Tattooists and body piercers currently undertake unaccredited courses, often run by private companies, and/or undergo local training and assessment against locally determined competencies through self-funded apprentice/mentorship schemes. These training methods vary in terms of quality, duration and content.

The absence of tattoo and body piercing accredited training and competencies is an area that needs to be addressed nationally and is outside the scope of this guidance. From an infection control viewpoint, any agreed national training and competencies for tattooists and body piercers should include the infection control areas listed in the procedure manuals/policies section (below).

#### **PROCEDURE MANUALS/POLICIES**

It is recommended that practitioners produce a written procedure/policy manual for use by staff. It should be based on evidence – based guidance and be easily available and be easily understood by all groups of staff. Policies/procedures should be reviewed annually so that practice is up to date and should indicate ownership (i.e. who is responsible for managing the policy) and authorship.

Policies/procedures should include:

- Hand washing procedure.
- Cleaning policy and rota.
- Decontamination procedures.
- Management of waste.
- Management of blood spillages.
- Use of personal protective equipment (PPE).

- Needle stick injury (and basic first aid practice in relation to this).
- Safe sharps handling/disposal.
- COSHH risk assessment/safe handling of chemicals.
- Training/education of staff.
- Staff health including hepatitis B vaccination status.

#### **AUDIT AND QUALITY MONITORING**

Services, policies and practices should be monitored on a regular basis by the premises manager/lead practitioner, not only to ensure practice is up to date and evidence based, but for quality purposes as well. By having written, up-to-date policies and procedures, documentation of staff training and proof of written, evidence based procedures/policies being followed, a premises can provide evidence of quality standards being maintained. This is useful when premises are asked by other professionals, and clients, for proof of good practice procedures and help demonstrate competence and quality in the event of a complaint.

Undertaking audits is part of providing evidence of quality performance. The template infection control audit tool which is included with this guidance could be used for undertaking infection control audits at a tattooing /body piercing premises. It provides a spreadsheet which gives percentage scores against good practice target scores in each section and overall.

It is suggested that audits are initially undertaken every six months (or more frequently if scores are below standard targets) and an action plan developed to address any gaps in audit findings. As a minimum an audit should be undertaken annually. It is important that there are named individuals designated against each action and that a deadline is given for each action so that progress can be identified against these by designated staff, as well allowing monitoring of progress to be followed by the manager/owner. A repeat audit allows the recording of changes made, as well as good practice to be monitored over time.

# Tattooing and body piercing guidance

## PART A

### Section 07 Governance

Click on text to view

**TRAINING AND COMPETENCIES**

**PROCEDURE MANUALS/ POLICIES**

**AUDIT AND QUALITY MONITORING**

**RECORD KEEPING**

**CONSENT AND AFTERCARE DOCUMENTATION**

See Appendix 08– contains an example of a consent form and there are a variety of written aftercare leaflets available

See Appendix 09– Aftercare follow-up record sheet

**References**

#### RECORD KEEPING

Accurate records are invaluable if infection problems occur and may assist the practitioner when investigations are conducted – for example, for verifying procedures performed and equipment check-tests carried out, when they were performed and on whom/by whom. It is important to keep accurate records of every client including:

- Full name, address, telephone number, date of birth and proof of age if needed.
- Relevant medical history/ allergies.
- Consent signature of client/ parent.
- Date and type of procedure conducted, site of procedure, type of jewellery (if applicable).
- The name of the practitioner.

All records should be used safely and stored securely, maintaining client confidentiality (e.g. locked paper records, safe use of computers to ensure clients details are not accessible by the general public or others with no legitimate reason to access them. Records should be kept safely on the premises named in the licence for a period of no less than 3 years (Dartford Borough Council 2009).

Staff training records should also be kept on site, as well as health and safety records such as risk assessments, an accident/incident book, and a log book with details of regular equipment checks.

#### CONSENT AND AFTERCARE DOCUMENTATION

Consent forms must be signed before any procedure is commenced. Both written and verbal aftercare information should be provided to the client as evidence of good and safe practice.

**See Appendix 08 – contains an example of a consent form and there are a variety of written aftercare leaflets available**

AFTERCARE FOLLOW-UP RECORD SHEET <small>(ATTACH TO CONSENT FORM)</small>		
<b>NAME OF CLIENT:</b> _____		
Date & time	Description of any concern or problem Action taken/advice given	Signature of practitioner

**See Appendix 09 – Aftercare follow-up record sheet**

#### References

Dartford Borough Council (2009). Code of practice for hygienic piercing. ([www.dartford.gov.uk](http://www.dartford.gov.uk))

## Section 08

Click on text to view

**DEFINITION OF AN OUTBREAK OR INCIDENT**

**OUTBREAK/INCIDENT INITIAL RESPONSE**

**References**

# Tattooing and body piercing guidance

## PART A

### Management of infectious disease incidents relating to tattooing and skin piercing

#### DEFINITION OF AN OUTBREAK OR INCIDENT

An infectious disease related outbreak or incident can be defined as:

- An incident in which two or more people experiencing a similar illness are linked in time/place.
- A greater than expected rate of infection compared with the usual background rate for the place and time where the outbreak has occurred.

#### OUTBREAK/INCIDENT INITIAL RESPONSE

In Northern Ireland, incidents/outbreaks are recognised using surveillance methodology by the Health Protection Team within the Public Health Agency, local authorities or microbiologists in regional laboratories. As soon as it becomes apparent that an incident/outbreak may exist, immediate contact between these parties is essential.

A risk assessment is undertaken following receipt of initial information and a decision made as to whether an outbreak or incident exists. In order to inform all subsequent decisions and actions, key facts would be established by the duty (or on-call) Health Protection Team.

An outbreak/incident is usually declared by the Consultant in Communicable Disease Control (CCDC) or Consultant in Health Protection (CHP) after consultation with a consultant microbiologist and/or senior environmental health officer. The involvement and assistance of the tattoo/body piercing premises concerned would be a high priority.

Once an outbreak/incident is declared, a multi-agency outbreak/incident control team would be set up to fully investigate the incident, ensure control measures are in place and a report generated with lessons learned. All legal powers relating to the investigation of outbreaks lie with the local authority. An outbreak/incident control team would give due consideration to the possibility of legal

proceedings, and if required seek guidance regarding the chain of evidence for a potential prosecution.

In the case of a tattooing/body piercing setting, most incidents would be expected to relate to suspected blood borne virus transmission to another person (be that from a practitioner to a client, from a client to the practitioner or from client to client via contaminated equipment or materials).

The infection control practices in place within the establishment would be reviewed as a key part of any incident investigation.

#### References

Health Protection Agency (2011). London Infectious Disease Outbreak Management Plan.

Health Protection Agency (2012). The Communicable Disease Outbreak Plan V1.3. (Internal document).

## Tattooing and body piercing guidance

### PART B – Appendix 01

#### Infection, its causes and spread (continued including a glossary of infection-related terms)

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

#### THE CAUSES OF INFECTION

Numerous microorganisms harmlessly colonise the skin and the mucosal surfaces to form the normal flora of the human body. The presence of microorganisms does not constitute an infection. Colonising microorganisms cause no damage and often provide benefit to the person. It is when there is associated tissue damage that an infectious disease exists. Potential pathogens can also act as colonisers such as *Staphylococcus aureus*

#### GLOSSARY OF INFECTION-RELATED TERMS:

**A pathogen** is an organism capable of invading the body and causing disease. Such an organism is termed pathogenic (Bannister et al, 2006)

**An infectious disease** is an illness caused by a pathogen, which invades body tissues and causes damage. Not all infectious diseases spread from person to person, e.g. Legionnaires’ disease.

**A communicable disease** is an infectious disease that is capable of spreading from person to person, e.g. measles, tuberculosis.

#### Self infection (endogenous infection)

An infection that arises from the person’s own body flora e.g. bacteria that colonise the skin get into a break in the skin (wound) and cause an infection such as an abscess caused by *Staphylococcus aureus*.

#### Cross infection (exogenous infection)

This is an infection that arises from an external source e.g. from another person or via the environment.

#### GROUPS OF ORGANISMS CAPABLE OF CAUSING INFECTION.

Pathogens relevant to body art can be classified into:

**Bacteria** are single celled organisms of approximately one-thousandth to five-thousandth of a millimetre in diameter.

Bacteria can replicate independently and some bacteria can form spores that survive in the environment for long periods of time, e.g. *Mycobacterium tuberculosis*, Group A Streptococcus, *Salmonella Enteritidis*. Antibiotics are used to treat bacterial infections; bacteria can develop resistance to antibiotics, e.g. MRSA (meticillin resistant *Staphylococcus aureus*).

**Viruses** are smaller than bacteria and cannot replicate independently but grow inside the host’s cells. Viruses cannot be treated with antibiotics; there are a few anti-viral drugs available that are active against a limited number of viruses such as influenza. Many common viral infections resolve without treatment, e.g. measles, mumps, and rubella.

**Pathogenic Fungi** can be either moulds or yeasts. Infections caused by moulds or yeasts include ringworm caused by *Trichophyton rubrum* and thrush, which is a common yeast infection caused by *Candida albicans*.

#### TRANSMISSION (SPREAD) OF INFECTION

How an infection is spread (transmitted) varies according to the type of microorganism. Some microorganisms may be transmitted by more than one route:

- Direct or indirect contact, e.g. *Herpes simplex* virus, respiratory syncytial virus, *Staphylococcus aureus*.
- Respiratory droplets, e.g. influenza virus, mumps, *Bordetella pertussis* (whooping cough).
- Airborne, e.g. pulmonary tuberculosis, measles, chickenpox.
- Other infectious agents, such as blood-borne viruses, e.g. hepatitis B and C and D viruses (HBV, HCV, HDV) and HIV are transmitted rarely in healthcare settings, via percutaneous (sharps/ needles) or mucous membrane exposure (blood/ body fluid splashes to eyes/ mouth/ open wounds).



# Tattooing and body piercing guidance

## PART B – Appendix 02

### Infection, its causes and spread (including a glossary of infection-related terms)

#### Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

#### **IN TATTOOING AND SKIN PIERCING PROCEDURES, INFECTION TRANSMISSION CAN OCCUR BY DIRECT CONTACT AND INDIRECT CONTACT.**

##### **Direct contact (person to person)**

Direct transmission occurs when microorganisms are transferred from one infected person to another person without a contaminated intermediate object or person. Examples of direct contact include:

- Blood or other body fluids (including blood stained) that enter the body through contact with a mucous membrane or breaks (i.e. piercings, cuts, abrasions) in the skin.

##### **Indirect contact**

Indirect spread of infection is said to occur when an intermediate carrier is involved in the spread of pathogens such as hands, fomites or vectors.

- **Hands** – The hands of the practitioners are probably the most important vehicles of cross-infection within the tattooing and skin piercing environment. The hands of staff and clients can carry microbes to other body sites, equipment and staff. Therefore, promotion of hygienic practices for everyone is the key to preventing and controlling infections.
- A **fomite** is defined as an object that becomes contaminated with infected organisms and which subsequently transmits those organisms to another person. Examples of potential fomites are instruments or practically any inanimate article e.g. contaminated needles/ tattooing equipment (blood-borne viruses).

##### **References**

Bannister B, Gillespie S and Jones J (2006). Infection Microbiology and Management. 3rd edition. Blackwell Publishing.

# Tattooing and body piercing guidance

## PART B – Appendix 02

### Blood borne viruses

#### Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

Blood borne viruses (BBVs) are viral infections which are spread through infected blood and body fluids, such as semen. The BBVs of greatest concern in the tattoo and piercing industry are hepatitis B virus (HBV), hepatitis C virus (HCV), hepatitis D virus (HDV) and human immunodeficiency virus (HIV). Hepatitis B and C viruses infect and damage the liver. Hepatitis D (HDV) requires the presence of the hepatitis B virus to survive in the body. This means that it is only possible to have hepatitis D if one also has hepatitis B. HIV stops a person's body from fighting infections properly.

BBVs can be passed on or 'transmitted' if an infected person's blood is able to enter another person's bloodstream. This can happen in a number of different ways:

- Sexual contact, both heterosexual and homosexual.
- Infected blood passing from one person to another e.g. through cuts or damaged skin.
- Sharing razors or toothbrushes.
- Sharing needles and syringes.
- Through ear piercing and other types of body piercing, tattooing and acupuncture if equipment is not properly sterilized.
- From mother to baby during or after pregnancy.
- All blood donations are now screened for hepatitis B virus, but before this started it was possible to become infected through blood transfusions.

The infection is not passed on through everyday activities such as coughing, sneezing, shaking hands or sharing food, crockery, baths or toilets.

BBVs can cause serious, chronic diseases (e.g. liver cirrhosis, cancer) or even death to the individual affected. If an individual has an ongoing (chronic) infection, they will pose a continuing risk of infection to others.

Sometimes an individual may be infected with a blood borne virus, but not be aware that they have the infection and that they are therefore an infectious risk to others.

Infections from BBVs can be prevented or avoided in the tattoo and body piercing setting, if robust infection control practices (including immunisation against hepatitis B) are used by all practitioners at all times when dealing with anyone else's blood and body fluids.

#### Reference:

British Liver Trust (<http://www.britishlivertrust.org.uk>)

# Tattooing and body piercing guidance

PART B – Appendix 03  [Click to download PDF version](#)

## Safe use and disposal of sharps

### Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

[↪ Main contents](#)

### Before use:

- Ensure that sharps disposal box is correctly assembled.
- Ensure that the label on the box is filled in upon assembly.
- Ensure appropriate colour sharps box lid for use based on medicinal contamination and how the waste should be treated and disposed of i.e. orange lid for sharps derived from tattooing/body piercing procedures.
- Sharps boxes are type approved for solids and should not be used for quantities of liquid waste.
- Sharps boxes must comply with UN 3291 and BS7320 standards.
- Boxes must be available in different sizes. Tamper-proof sharps containers are also available.
- Boxes must be available at all locations where sharps are used.
- Boxes must never be placed on the floor.
- Boxes must be placed on a level surface or wall-mounted below shoulder height and be near to the area they are being used.
- Boxes must never be left in areas where clients may have open access to them.
- Assess, in terms of risk, the most appropriate size of sharps container for the tattoo/body piercing setting.

### During use:

- Practitioners must be competent in procedures using sharps.
- The person using the sharp is responsible for disposing of it.
- Never pass sharps from hand to hand.
- Wear appropriate personal protective equipment (gloves at a minimum).
- Assemble devices with care.
- Do not disassemble devices (e.g. needle bar and needle)—dispose of as a complete unit.
- Do not re-sheath/recap used needles/razors.
- Close sharps box opening (temporary closure device) between uses.
- Never move an open sharps box.
- Use the handle to carry.

### After use:

- Disposal of sharps is the responsibility of the user.
- Dispose of sharps immediately after use.
- Do not bend or break needles before disposal (e.g. a tattoo needle from a needle bar).
- Do not leave full sharps boxes for disposal by other staff.
- Fill sharps boxes only to the 'fill' line and never overfill.
- Shut and lock box when full for disposal.
- Never use tape to seal sharps boxes.
- Label box with source such as name of person/ premises and describe waste content.
- Dispose of sharps boxes as clinical waste for incineration only.
- Never place sharps boxes in clinical/offensive waste bags.
- Sharps containers must never be left unsupervised. They must be locked in a cupboard/ operating/procedure room when not in use.
- Never try and retrieve items from a sharps container.
- Place damaged sharps containers inside a larger container – lock and label prior to disposal. Do not place inside a waste bag.

### Reference

National Institute for Clinical Excellence (NICE) (2012). Infection control: Prevention and control of healthcare-associated infections in primary and community care (CG139).

<http://guidance.nice.org.uk/CG139>

## Tattooing and body piercing guidance

PART B – Appendix 04  Click to download PDF version

### First Aid following a blood/body fluid exposure

Click on text to view

Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms

Appendix 02 – Blood borne viruses

Appendix 03– Safe use and disposal of sharps

Appendix 04– First Aid following a blood/body fluid exposure

Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill

Appendix 06 – Principles for good waste handling

Appendix 07– Template protocol for environmental cleaning of premises

Appendix 08– Tattooing/body piercing consent form

Appendix 09– Aftercare follow-up record sheet


Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing

Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners

Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers

Appendix 13– Autoclave daily record sheet

↪ Main contents



**The priority is to seek advice/ medical attention immediately, at a local Emergency Department, ideally within one hour of the injury occurring**

Reference:  
Health and Safety Executive (2012) How to deal with an exposure incident. Blood Borne Viruses Guidance. (<http://www.hse.gov.uk/biosafety/blood-borne-viruses/how-deal-exposure-incident.htm>)

## Tattooing and body piercing guidance

PART B – Appendix 05  Click to download PDF version

### Protocol for cleaning up blood or a blood stained body fluid spill

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

#### CLEANING UP BLOOD OR A BLOOD STAINED BODY FLUID SPILL

- Prevent access to the area containing the spillage until it has been safely dealt with.
- Obtain chlorine based spill kit.
- Put on apron and gloves
- Apply disinfectant granules to the spill, this congeals the spill to enable easier cleaning of the area OR make up and use the disinfectant solution as per product instructions to a dilution of 10,000ppm available chlorine. Leave in place for the designated exposure time (at least two minutes). Ensure the surface can tolerate chlorine.

#### DO NOT USE MOPS TO CLEAN UP BLOOD.

- Use the scoop and scraper (or disposable paper towels) to pick up the congealed body fluid and place in the appropriate waste bag.
- Using the disposable paper towels and disinfectant solution clean area thoroughly and dry afterwards. Detergent and warm water can also be used afterwards.
- Ensure all equipment used is disposed of in the appropriate waste bag and then finally remove gloves and apron and place in the waste bag.
- Wash hands.
- Ensure that the waste bag is placed in the appropriate disposal bin/container immediately after use.
- Damp-mop the affected area.

#### MANAGING BLOOD SPOTS

Apply chlorine based disinfectant solution to a wet paper towel and clean spillage area. Discard waste as above.

#### IF BLOOD/BODY FLUID SPLASHES INTO THE EYES OR MOUTH

Rinse freely with water. Seek immediate medical advice if the splash gets into the mucous membrane.

#### BLOOD SPILLS ON CLOTHING

Change clothes (immediately if possible) and place into a plastic bag. Wash clothes as soon as possible in a hot cycle.

#### CLEANING UP VOMIT OR URINE SPILLS

Chlorine-based disinfectants will give off highly toxic gas if mixed with acidic substances. Ideally chlorine-based products should not be used on vomit and urine due to the slight risk of chlorine gas being released. To clean up vomit or urine spills, follow the same process as above but replace chlorine-based disinfectant granules with a non-chlorine based product, or use paper towels to absorb as much of the spillage as possible. Always clean areas with detergent and warm water. A chlorine-based disinfectant can be used to disinfect the area but only after the urine/vomit has been cleaned up.

#### IMPORTANT NOTE

**Chlorine-based disinfectants/ absorbent granules such as sodium dichloisocyanurate (NaDCC) should not be used on urine or vomit spills. NEVER mix chlorine-based disinfectants with any other cleaning/disinfectants. Hypochlorite solutions may bleach fabrics and other materials, as well as corrode metals, so care is needed regarding which surfaces they can be used upon.**

#### Reference

National Patient Safety Agency (2009). The Revised Healthcare Cleaning Manual. (<http://www.nrls.npsa.nhs.uk/resources/?EntryId45=61830>)



# Tattooing and body piercing guidance

PART B – Appendix 06  [Click to download PDF version](#)

## Principles for good waste handling

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

### GOOD WASTE HANDLING PRINCIPLES:

- Waste should be segregated at the point of origin.
- Bags/bins should only be filled to  $\frac{3}{4}$  full.
- Waste bags should be used in foot operated pedal bins.
- Waste bags should be sealed securely and marked with 'point of origin' label prior to disposal.
- Waste should be collected on a regular basis by a licensed waste management contractor.
- Personal protective clothing should be worn when handling waste.
- Waste should be correctly bagged in appropriate colour-coded bags which must be UN-approved and comply with BS EN ISO 7765:2004 and BS EN ISO 6383:2004.
- Waste should be double bagged where the exterior of the bag is contaminated or the original bag is split, damaged or leaking.
- Waste should be kept in a rigid-sided, fire retardant holder or container with a foot operated lid, and, so far as is reasonably practicable, out of the reach of children and unauthorised personnel.
- Waste should be stored in a labelled, lockable/secure, vermin-proof storage space for collection, on a well-drained, impervious hard standing floor, which is provided with wash-down facilities.
- Bags should be securely sealed and labelled with coded tags at the point of use to identify their source.
- Waste should not be decanted into other bags, regardless of volume; be contaminated on the outside or re-used.
- 'Sharps' must be disposed of into approved sharps containers that meet BS 7320/UN 3291.
- 'Sharps' containers should **NEVER** be placed into any waste bag.

## Tattooing and body piercing guidance

PART B – Appendix 07  [Click to download PDF version](#)

### Template protocol for environmental cleaning of premises

**Click on text to view**

**Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms**

**Appendix 02 – Blood borne viruses**

**Appendix 03– Safe use and disposal of sharps**

**Appendix 04– First Aid following a blood/body fluid exposure**

**Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill**

**Appendix 06 – Principles for good waste handling**

**Appendix 07– Template protocol for environmental cleaning of premises**

**Appendix 08– Tattooing/body piercing consent form**

**Appendix 09– Aftercare follow-up record sheet**

**Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing**

**Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners**

**Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers**

**Appendix 13– Autoclave daily record sheet**

ITEM	FREQUENCY	METHOD
High risk treatment surfaces	After use	Treatment area surfaces cleaned and dried between clients using detergent and then disinfected using a bleach solution (1000 ppm)  Use disposable cloths/paper towels
Non high risk surfaces	At least daily	Use general-purpose detergent  Dry thoroughly  Use disposable cloths/paper towels
Hand wash basins and sinks	Daily	Standard detergent
Floors	Daily	Mop with water and detergent  Disinfectant is required only after contamination with blood spillages
Bins	As required	Empty bins daily.  If contaminated, clean with water and detergent and then disinfect
Couches	Between clients	Wipe with hot, soapy water and dry thoroughly  Clean with disinfectant against blood borne viruses if contaminated with blood
Walls/ceilings	As required	Routine cleaning not required  Clean periodically with water and general purpose detergent  Clean with disinfectant against blood borne viruses if contaminated with blood

Adapted from National Patient Safety Agency (2009). The Revised Healthcare Cleaning Manual. (<http://www.nrls.npsa.nhs.uk/resources/?EntryId45=61830>)

## Tattooing and body piercing guidance

### PART B – Appendix 08

#### Tattooing/body piercing consent form

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

 Click to download Word document version

 Click to download PDF version

#### Tattooing/Body Piercing Consent Form

Name of Premises:	
Address & Tel No of Premises:	
Name of Practitioner (print):	
Name of Client (print):	
Address & Tel No of client:	
Age of Client and DOB:	
Age ID of Client Seen:	Yes / No
Type of Procedure:	Tattoo / Body Piercing Description:
Site of Procedure and design if applicable):	
Type of Jewellery Used (as applicable):	

# Tattooing and body piercing guidance

## PART B – Appendix 09

### Aftercare follow-up record sheet

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)



Click to download  
Word document version



Click to download  
PDF version

#### AFTERCARE FOLLOW-UP RECORD SHEET (ATTACH TO CONSENT FORM)

**NAME OF CLIENT:**

Date & time	Description of any concern or problem Action taken/advice given	Signature of practitioner

## Tattooing and body piercing guidance

PART B – Appendix 10  [Click to download PDF version](#)

### Decontamination requirements for equipment used in tattooing and skin piercing

[Click on text to view](#)

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

[↩ Main contents](#)

Equipment for	Application of item	Recommendations
<b>Tattooing:</b>		
Holders for needles i.e. tube, tip and grip	Hold needles that pierce skin	Dismantle then clean and sterilize, or single-use
Needles and needle bars	Pierce skin	Single-use pre-sterilized
Ink caps	Hold inks that will become contaminated with blood	Single-use
Ink cap trays	Hold ink caps and will become contaminated with blood during use	Either single-use or clean and sterilize
Motors & clipcords	Will become contaminated with blood via the practitioner’s hands	Reduce contamination by covering with impervious barrier (e.g. plastic sleeving)  Remove impervious barrier carefully after use minimising transfer of contamination to the item  Thoroughly remove any visible contamination with detergent, then dry  Disinfect surface with 70% alcohol
Elastic bands	Will become contaminated with blood via the practitioner’s hands	Single-use
<b>Body piercing:</b>		
Needles, cannulas	Pierce skin	Single use pre-sterilized
Clamps used for skin folds, looped forceps, pliers and receiving tubes	In close contact with pierced skin	Clean and sterilize or single-use
Jewellery	Inserted into sterile body tissues	Sterilize or pre-sterilized
Spatulas, cotton/gauze pads and paper towels	Will become contaminated with blood	Single-use
Vernier calipers	Should only be used on clean, unbroken skin	Clean between uses  Single use or autoclavable alternative should be used for oral and genital piercings
Ear piercing guns	These should only be of the type that use single-use self-contained cartridge containing the stud and back, such that the gun itself makes no client contact	Clean according to manufacturer’s instructions
Cartridges used with ear piercing guns		Single-use



## Tattooing and body piercing guidance

PART B – Appendix 11 [Click to download PDF version](#)

Equipment sterilization standard – self assessment and decision making tool for tattoo and body piercing practitioners

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)


[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)



## Tattooing and body piercing guidance

PART B – Appendix 12 [Click to download PDF version](#)

Equipment and body piercing jewellery sterilization standard for tattooists and body piercers

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)


[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)



## Tattooing and body piercing guidance

### PART B – Appendix 13

#### Autoclave daily record sheet

Click on text to view

[Appendix 01 – Infection, its causes and spread, including a glossary of infection-related terms](#)

[Appendix 02 – Blood borne viruses](#)

[Appendix 03– Safe use and disposal of sharps](#)

[Appendix 04– First Aid following a blood/body fluid exposure](#)

[Appendix 05– Protocol for cleaning up a blood or a blood stained body fluid spill](#)

[Appendix 06 – Principles for good waste handling](#)

[Appendix 07– Template protocol for environmental cleaning of premises](#)

[Appendix 08– Tattooing/body piercing consent form](#)

[Appendix 09– Aftercare follow-up record sheet](#)

[Appendix 10 – Decontamination requirements for equipment used in tattooing and skin piercing](#)

[Appendix 11– Equipment sterilization standard- self assessment and decision making tool for tattoo and body piercing practitioners](#)

[Appendix 12 – Equipment and body piercing jewellery sterilization standard for tattooists and body piercers](#)

[Appendix 13– Autoclave daily record sheet](#)

 Click to download Word document version

 Click to download PDF version

#### AUTOCLAVE DAILY RECORD SHEET

Please keep these records in date order for inspection

Autoclave Type	Serial Number
Week Commencing	Location
Type of Water used (ideally sterile water for irrigation)	

Daily test	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Cycle Counter Number	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Time to reach holding temp	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Temp during holding period	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Pressure during holding period	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Total time at holding temp/pressure	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Water drained at end of day where appropriate	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Process check used	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Printout attached	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Initials of authorised user	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Weekly Safety Test	Yes/No	Comments
Door seals secure		
Door safety devices functioning correctly		
Safety Valves operating correctly		
Yearly service by a competent engineer		
Comments		
Name	Date	Signature

PLEASE KEEP THESE RECORDS IN A RING BINDER FOR SELF AUDIT/INSPECTION

# Tattooing and body piercing guidance

## PART C – Leaflet 01

### Tattoo aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

[Leaflet 04 – Body and surface piercing aftercare](#)

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Tattoo *aftercare*

### Key Advice

The aftercare following a tattoo is important to promote good healing and prevent the risk of infection.

For the first week or so it is normal for the area to be red and tender.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare.

The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the tattoo
- Keeping the tattooed area covered with non-stick gauze which is secured with hypo-allergenic tape.

### Hand washing

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

### Tattoo aftercare

Good practice is to cover the tattooed area with non-stick gauze which is then secured with hypo-allergenic tape. Gauze permits ventilation and aids healing.

A tattoo covering a large area may need to be covered with a sterile, non-adhesive dressing, at least during your journey home. However, simply keeping the area clean and dry is likely to be the best approach.

A tattoo covering a large area may have plastic film wrap applied, this must be clean (taken straight from the pack and used immediately) and you should be advised when to replace this covering by your practitioner.

Any cream that you apply must be used from an appropriate pot/tube at home and you should wash your hands before application. Cream can be purchased from your practitioner or a pharmacist.


Antibiotic creams should not be used except if infection has occurred and under supervision of your Doctor.

### Signs of infection

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your tattoo or if there are any signs of an allergic reaction to any of the products used.







Your practitioner is:

June 2013

**For further advice or information:**  
Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by:  
Public Health England North West  
Public Health England East Midlands  
Tattoo and Piercing Industry Union

# Tattooing and body piercing guidance

## PART C – Leaflet 02

### Ear and face piercing aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

[Leaflet 04 – Body and surface piercing aftercare](#)

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Ear and face piercing *aftercare*

**Key Advice**

The aftercare of body piercing is important to promote good healing and prevent the risk of infection.

Healing times for piercing will vary with the type and position of the piercing and vary from person to person.

For the first few weeks it is normal for the area to be red, tender and swollen.

Approximate healing times for the various piercings are:

Ear lobe piercing – 6 to 8 weeks  
 Ear cartilage piercing – 6 to 8 weeks  
 Cheek piercing – 2 to 3 months  
 Eyebrow piercing – 2 to 4 months  
 Nose piercing – up to 6 months

Remember these times are approximate and will depend on how healthy you are and whether you look after the piercing properly until healed.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare.

The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the piercing.
- Keeping the piercing clean.

**Hand washing**

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

A new piercing can be tender, itchy and slightly red and can remain so for a few weeks. A pale, odourless fluid may sometimes discharge from the piercing and form a crust. This should not be confused with pus, which would indicate infection.

**Ear piercing aftercare**

Including Lobes/ Tragus/ Anti Tragus/ Conch/ Helix/ Snug/ Diath/ Industrial/ Rook/ Translobal/ Transverse Lobe.

**Facial piercing aftercare**

Including Eyebrow, Bridge, Jestum, Vertical Labret, Septum/ Nostril.

Soak the piercing for a few minutes by submerging the area of skin containing the piercing in a clean jug or bowl containing a warm water solution (1/4 level teaspoon of preferably sea salt to an egg cup/shot glass of warm water). Alternatively wet a clean cloth or gauze in the solution and apply as a warm compress. This will soften any discharge and allow you to clean the entry and exit points of the piercing with a cotton bud or gauze. Once the discharge is removed or softened then jewellery can be gently moved so as to work a little warm water through the piercing. When cleaning always tighten the ball on any bars by screwing the ball to the right.

Do this twice each day, preferably after washing or bathing.

You can also use mild antibacterial solutions and soaps to wash the wound site of an ear piercing. Ask your local pharmacist to advise you and always follow the manufacturers' instructions. If irritation, redness or drying occurs discontinue use. Antibacterial wash is NOT suitable for nostrils, septum or vertical lips due to the tissue's delicate nature.

Dry the piercing using ONLY fresh disposable paper towel/kitchen roll. A communal hand/bath towel should never be used.

**Note:** Cartilage piercings occasionally form lumps commonly known as granulomas. This is just trapped fluid and can easily be resolved using the heat and pressure from a warm water compress once a day, replacing one of your daily cleanings.

**Expect** some swelling and soreness from your new piercing. Any knock or bang can cause swelling or soreness to flare up again throughout the healing phase.

**Do not** use cotton wool to clean the piercing as the fibres in the cotton wool may get caught in the piercing.

**Do not** pick at any discharge and **do not** move, twist or turn the piercing whilst dry. If any secreted discharge has hardened then turning jewellery may cause the discharge to tear the piercing, allowing bacteria to enter the wound and prolonging the healing time.

**Do not** use sunbeds for the first two weeks, or if you decide to then cover the wound area with a breathable plaster during tanning.

**Do not** swim for the first 24 hours following a piercing.

**Signs of infection**

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your piercing or if there are any signs of an allergic reaction to any of the products used.

**For further advice or information:**

Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by: Public Health England North West Public Health England East Midlands Tattoo and Piercing Industry Union

Your practitioner is:

June 2013



# Tattooing and body piercing guidance

## PART C – Leaflet 03

### Oral piercing aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

**[Leaflet 03 – Oral piercing aftercare](#)**

[Leaflet 04 – Body and surface piercing aftercare](#)

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Oral piercing *aftercare*

**Key Advice**

The aftercare of body piercing is important to promote good healing and prevent the risk of infection.

Healing times for piercing will vary with the type and position of the piercing and vary from person to person.

For the first few weeks it is normal for the area to be red, tender and swollen.

An approximate healing times for oral piercing are:

Tongue piercing – 2 to 4 weeks.  
Lip – 3 to 6 weeks  
Cheek – 2 to 3 months

Remember these times are approximate and will depend on how healthy you are and whether you look after the piercing properly until healed.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare.

The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the piercing
- Keeping the piercing clean.

**Hand washing**

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

A new piercing can be tender, itchy and slightly red and can remain so for a few weeks. A pale, odourless fluid may sometimes discharge from the piercing and form a crust. This should not be confused with pus, which would indicate infection.

**Oral piercing aftercare**

For the internal healing of oral piercing including all piercing of the tongue, lip and cheek

Gargle after each meal with an alcohol-free mouthwash or a warm salt water solution (1/4 level teaspoon of preferably sea salt to an egg cup/shot glass of warm water).

For the external healing of oral piercing: including all piercings to the lip and cheek

Wet a clean cloth or gauze in the warm salt water solution and apply as a warm compress. This will soften any discharge and allow you to clean the entry and exit points of the piercing with a cotton bud or gauze dipped into the warm salt water solution. Once the discharge is removed or softened then jewellery can be gently moved so as to work a little warm water through the piercing. When cleaning always tighten the ball on any bars by screwing the ball to the right.

Do this twice each day, preferably after washing or bathing.

Dry the piercing using ONLY fresh disposable paper towel/kitchen roll. A communal hand/bath towel should never be used.

**Do not** use cotton wool to clean the piercing as the fibres in the cotton wool may get caught in the piercing.

Your practitioner is:

**Do not** pick at any discharge and **do not** move, twist or turn the piercing whilst dry. If any secreted discharge has hardened then turning jewellery may cause the discharge to tear the piercing, allowing bacteria to enter the wound and prolonging the healing time.

**Do not** use sunbeds for the first two weeks, or if you decide to then cover the wound area with a breathable plaster during tanning.

**Do not** swim for the first 24 hours following a piercing.

**Special aftercare for tongue piercing**

For the first few days take care when eating and avoid spicy foods.

Cold products such as ice and ice cream can help reduce swelling.

Refrain from oral sex of any description until the piercing has fully healed.

**Signs of infection**

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your piercing or if there are any signs of an allergic reaction to any of the products used.

**For further advice or information:**

Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by: Public Health England North West Public Health England East Midlands Tattoo and Piercing Industry Union






June 2013

# Tattooing and body piercing guidance

## PART C – Leaflet 04

### Body and surface piercing aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

**[Leaflet 04 – Body and surface piercing aftercare](#)**

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Body and surface piercing *aftercare*

**Key Advice**

The aftercare of body piercing is important to promote good healing and prevent the risk of infection.

Healing times for piercing will vary with the type and position of the piercing and vary from person to person.

For the first few weeks it is normal for the area to be red, tender and swollen.

Approximate healing times for surface, navel and nipple piercing can be as long as 6 months to 1 year.

Remember these times are approximate and will depend on how healthy you are and whether you look after the piercing properly until healed.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare. The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the piercing
- Keeping the piercing clean.

**Hand washing**

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

A new piercing can be tender, itchy and slightly red and can remain so for a few weeks. A pale, odourless fluid may sometimes discharge from the piercing and form a crust. This should not be confused with pus, which would indicate infection.

**Body and surface piercing aftercare** including Nape, Horizontal Navel, Nipple, Horizontal Eyebrow, Wrist, Anti-eyebrow, Madison, Chin, Vertical Bridge, Pubic, Handweb and Prayer.

Soak the piercing for a few minutes by submerging the area of skin containing the piercing in a clean jug or bowl containing a warm water solution (1/4 level teaspoon of preferably sea salt to an egg cup/shot glass of warm water). Alternatively wet a clean cloth or gauze in the solution and apply as a warm compress. This will soften any discharge and allow you to clean the entry and exit points of the piercing with a cotton bud or gauze. Once the discharge is removed or softened then jewellery can be gently moved so as to work a little warm water through the piercing. When cleaning always tighten the ball on any bars by screwing the ball to the right.

Do this twice each day, preferably after washing or bathing.

You can also use mild antibacterial solutions and soaps to wash the wound site. Ask your local pharmacist to advise you and always follow the manufacturers' instructions. If irritation, redness or drying occurs discontinue use.

Dry the piercing using **ONLY** fresh disposable paper towel/kitchen roll. A communal hand/bath towel should never be used.

**Do not** use cotton wool to clean the piercing as the fibres in the cotton wool may get caught in the piercing.

Your practitioner is:

**Do not** pick at any discharge and **do not** move, twist or turn the piercing whilst dry. If any secreted discharge has hardened then turning jewellery may cause the discharge to tear the piercing, allowing bacteria to enter the wound and prolonging the healing time.

**Do not** wear tight clothing following nipple piercing

**Do not** wear tight clothing e.g. tights, belts or high waisted clothing after navel piercing as this may irritate the wound and delay healing.

**Do not** use sunbeds for the first two weeks, or if you decide to then cover the wound area with a breathable plaster during tanning.

**Do not** swim for the first 24 hours following a piercing.

**Signs of infection**

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your piercing or if there are any signs of an allergic reaction to any of the products used.

**For further advice or information:**  
Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by:  
Public Health England North West  
Public Health England East Midlands  
Tattoo and Piercing Industry Union





June 2013

# Tattooing and body piercing guidance

## PART C – Leaflet 05

### Genital piercing (female) aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

[Leaflet 04 – Body and surface piercing aftercare](#)

**[Leaflet 05 – Genital piercing \(female\) aftercare](#)**

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Genital piercing (female) *aftercare*

**Key Advice**

The aftercare of body piercing is important to promote good healing and prevent the risk of infection.

Healing times for piercing will vary with the type and position of the piercing and vary from person to person.

For the first few weeks it is normal for the area to be red, tender and swollen.

The healing time for a genital piercing can be from 2 to 12 weeks.

Remember these times are approximate and will depend on how healthy you are and whether you look after the piercing properly until healed.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare.

The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the piercing
- Keeping the piercing clean.

**Hand washing**

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

A new piercing can be tender, itchy and slightly red and can remain so for a few weeks. A pale, odourless fluid may sometimes discharge from the piercing and form a crust. This should not be confused with pus, which would indicate infection.

**Female Genital piercing aftercare** including Clitoral Hood, Inner and Outer Labia, Fouchette, Christina and Triangle.

Soak the piercing for a few minutes by submerging the area of skin containing the piercing in a clean container, such as a bowl containing a warm water solution (1/4 level teaspoon of preferably sea salt to an egg cup/shot glass of warm water). Alternatively wet a clean cloth or gauze in the solution and apply as a warm compress. This will soften any discharge and allow you to clean the entry and exit points of the piercing with a cotton bud or gauze. Once the discharge is removed or softened then jewellery can be gently moved so as to work a little warm water through the piercing. When cleaning always tighten the ball on any bars by screwing the ball to the right.

Do this twice each day, preferably after washing or bathing.

Dry the piercing using ONLY fresh disposable paper towel/kitchen roll. A communal hand/bath towel should never be used.

**Do not** use antibacterial products as they can kill the good bacteria that are naturally present.

**Do not** swim for the first 24 hours following a piercing.

**Do not** pick at any discharge and **do not** move, twist or turn the piercing whilst dry. If any secreted discharge has hardened then turning jewellery may cause the discharge to tear the piercing, allowing bacteria to enter the wound and prolonging the healing time.

**Refrain** from any type of sexual activity until the piercing has healed or is 'dry'.

**Always** use barrier protection such as condoms, otherwise you are at increased risk of acquiring a sexually transmitted infection.

**Signs of infection**

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your piercing or if there are any signs of an allergic reaction to any of the products used.



Your practitioner is:

June 2013

**For further advice or information:**  
 Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by:  
 Public Health England North West  
 Public Health England East Midlands  
 Tattoo and Piercing Industry Union






# Tattooing and body piercing guidance

## PART C – Leaflet 06

### Genital piercing (male) aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

[Leaflet 04 – Body and surface piercing aftercare](#)

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

**[Leaflet 06 – Genital piercing \(male\) aftercare](#)**

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Genital piercing (male) *aftercare*

**Key Advice**

The aftercare of body piercing is important to promote good healing and prevent the risk of infection.

Healing times for piercing will vary with the type and position of the piercing and vary from person to person.

For the first few weeks it is normal for the area to be red, tender and swollen.

The healing time for a genital piercing can be from 2 to 12 weeks.

Remember these times are approximate and will depend on how healthy you are and whether you look after the piercing properly until healed.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare.

The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the piercing
- Keeping the piercing clean.

**Hand washing**

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

A new piercing can be tender, itchy and slightly red and can remain so for a few weeks. A pale, odourless fluid may sometimes discharge from the piercing and form a crust. This should not be confused with pus, which would indicate infection.

**Male Genital piercing aftercare**

including glans penis piercing such as PA, Reverse PA, Apadroya, Ampallang, Dydoe and Frenum, as well as for other piercings including Hafada, Scrotum, Foreskin and Guiche.

Soak the piercing for a few minutes by submerging the area of skin containing the piercing in a clean container, such as a bowl containing a warm water solution (1/4 level teaspoon of preferably sea salt to an egg cup/shot glass of warm water). Alternatively wet a clean cloth or gauze in the solution and apply as a warm compress. This will soften any discharge and allow you to clean the entry and exit points of the piercing with a cotton bud or gauze. Once the discharge is removed or softened then jewellery can be gently moved so as to work a little warm water through the piercing. When cleaning always tighten the ball on any bars by screwing the ball to the right.

Do this twice each day, preferably after washing or bathing.

When cleaning always tighten the ball on any bars by screwing the ball to the right.

Dry the piercing using ONLY fresh disposable paper towel/kitchen roll. A communal hand/bath towel should never be used.

**Do not** use antibacterial products as they can kill the good bacteria that are naturally present.

**Do not** swim for the first 24 hours following a piercing.

**Do not** pick at any discharge and **do not** move, twist or turn the piercing whilst dry. If any secreted discharge has hardened then turning jewellery may cause the discharge to tear the piercing, allowing bacteria to enter the wound and prolonging the healing time.

**Refrain** from any type of sexual activity until the piercing has healed or is 'dry'.

**Always** use barrier protection such as condoms, otherwise you are at increased risk of acquiring a sexually transmitted infection.

**Signs of infection**

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your piercing or if there are any signs of an allergic reaction to any of the products used.

**For further advice or information:**

Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by: Public Health England North West Public Health England East Midlands Tattoo and Piercing Industry Union



Your practitioner is:

June 2013








# Tattooing and body piercing guidance

## PART C – Leaflet 07

### Microdermal implants aftercare

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

[Leaflet 04 – Body and surface piercing aftercare](#)

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)

[Poster – How to handwash](#)

Click to download PDF

## Microdermal implants *aftercare*

**Key Advice**

The aftercare of body piercing is important to promote good healing and prevent the risk of infection.

Microdermal implants or dermal anchors are small pieces of jewellery made from titanium inserted into the skin with the stem protruding above, onto which an attachment of your choice is fitted. This attachment should remain in place for at least three months or until the piercing has fully healed. Once it has fully healed other pieces of jewellery can be attached.

Healing times for piercing will vary with the type and position of the piercing and vary from person to person.

For the first few weeks it is normal for the area to be red, tender and swollen.

As with all body art, infection is a risk. To reduce these risks take advice from your practitioner regarding aftercare.

The risk of infection can be greatly reduced by good general hygiene including:

- Hand washing before touching the implant.
- Keeping the implant clean.

**Hand washing**

Hand washing is the single most important method of reducing infection. Hands must be washed prior to touching the affected area, therefore reducing the risk of infection.

Wash your hands in warm water and liquid soap, always dry your hands thoroughly with a clean towel or paper towel. This should remove most germs and prevent them being transferred to the affected area.

A new piercing can be tender, itchy and slightly red and can remain so for a few weeks. A pale, odourless fluid may sometimes discharge from the piercing and form a crust. This should not be confused with pus, which would indicate infection.

**Keeping the implant piercing clean**

The implant needs to be cleaned twice each day. Cleaning more frequently may damage the skin and slow down the healing process. Your practitioner may also advise you to soak the implant in warm salt water twice a week.

Make up a quantity of warm salt water solution (1/4 level teaspoon of preferably sea salt to an egg cup/shot glass of warm water).

Use a clean cloth or gauze dipped in the solution and apply as a warm compress and also to dab the area to make sure the area under the disc is cleaned as this may become encrusted.

This will soften any discharge and allow you to clean the piercing points with a cotton bud or gauze dipped into the warm salt water solution.

If the area around the implant becomes encrusted soak the piercing for a few minutes by submerging the area of skin containing the piercing in a clean jug or bowl containing the warm salt water solution and loosen the discharge using a cotton bud or clean floss.

Always dry the area thoroughly after cleaning your implant using ONLY fresh disposable paper towel/kitchen roll. A communal hand/bath towel should never be used.

**Do not** over clean the site as this may damage the skin around the implant

**Do not** change the cap of the microdermal implant until fully healed. If in any doubt take advice from your operator

**Do not** use cotton wool to clean the piercing as the fibres in the cotton wool may get caught in the piercing.

Your practitioner is:

**Do not** pick at any discharge and **do not** move, twist or turn the piercing whilst dry. If any secreted discharge has hardened then turning jewellery may cause the discharge to tear the piercing, allowing bacteria to enter the wound and prolonging the healing time.

**Do not** wear clothing that will rub against the piercing as this may irritate the wound and delay healing.

**Do not** use sunbeds for the first two weeks, or if you decide to then cover the wound area with a breathable plaster during tanning.

**Do not** swim for the first 24 hours following a piercing.

**Accidental damage or loss of disc**

Contact your practitioner if the implant gets caught in anything or the piercing becomes damaged.

In the unlikely event the disc breaks or comes off, return to the practitioner and have a new disc fitted immediately. If the disc is not replaced the implant may get lost under the skin and will require removal.

**Signs of infection**

If appropriate aftercare is not followed infection may occur. The signs of infection are:

- Swelling and redness that increases around the wound.
- A severe burning and throbbing sensation round the site.
- Increased tenderness and increasingly painful to touch.
- An unusual discharge (yellow or green) with an offensive smell.

Speak to your practitioner or seek medical attention immediately if you suffer from any of the above or have any concerns regarding infection in your piercing or if there are any signs of an allergic reaction to any of the products used.

**For further advice or information:**

Contact your local Environmental Health Department, or your local Public Health England Health Protection Team

This Information is provided by: Public Health England North West Public Health England East Midlands Tattoo and Piercing Industry Union



June 2013








## Tattooing and body piercing guidance

### PART C – Poster

#### How to handwash

Click on text to view

[Leaflet 01 – Tattoo aftercare](#)

[Leaflet 02 – Ear and face piercing aftercare](#)

[Leaflet 03 – Oral piercing aftercare](#)

[Leaflet 04 – Body and surface piercing aftercare](#)

[Leaflet 05 – Genital piercing \(female\) aftercare](#)

[Leaflet 06 – Genital piercing \(male\) aftercare](#)

[Leaflet 07 – Microdermal implants aftercare](#)


[Poster – How to handwash](#)

Click to download PDF

# How to Handwash?

WASH HANDS WHEN VISIBLY SOILED! OTHERWISE, USE HANDRUB

Duration of the entire procedure: 40-60 seconds




Wet hands with water;



Apply enough soap to cover all hand surfaces;



Rub hands palm to palm;



Right palm over left dorsum with interlaced fingers and vice versa;




Palm to palm with fingers interlaced;



Backs of fingers to opposing palms with fingers interlocked;




Rotational rubbing of left thumb clasped in right palm and vice versa;




Rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa;




Rinse hands with water;



Dry hands thoroughly with a single use towel;



Use towel to turn off faucet;



Your hands are now safe.



World Health Organization

Patient Safety  
A World Alliance for Safer Health Care

SAVE LIVES  
Clean Your Hands

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this document. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use. WHO acknowledges the Hôpitaux Universitaires de Genève (HUG), in particular the members of the Infection Control Programme, for their active participation in developing this material.

May 2009

# Tattooing and body piercing guidance

## PART D

### Audit tool

 [Click to download Excel document](#)

**INFECTION CONTROL AUDIT FOR TATTOO AND BODY PIERCING PREMISES\***

**NAME AND ADDRESS OF PREMISES:**  
**TELEPHONE NUMBER:**  
**OWNER/ LICENCEE'S NAME:**  
**OPERATOR'S NAME:**  
**EMAIL CONTACT DETAILS**

TYPE OF PREMISES (please tick as applicable)	
Tattooing	
Body Piercing	
Other (please state):	

**DATE OF AUDIT:**  
**NAME OF AUDITOR:**

<b>Indicators met:</b>	0
<b>Indicators not met:</b>	0
<b>Indicators not applicable:</b>	0
<b>Overall score:</b>	-


This audit tool is to be used by practitioners, primarily, to enable good practice to be captured and documented. It can also identify best practice gaps which will need action to be taken using an action planning process. It can be used as a discussion point between practitioner and local authority officers, as to what practice gaps need to be prioritised.

**Completion:**  
 Please allow approximately two hours to undertake the audit.  
 In the "Met" column, record "Yes", "No" or "N/A" against all indicators.

**To print all pages:**  
 Go to "print" and choose "entire workbook".

**Calculating scores:**  
**Automatic** - This tool will give you automated % scores for each standard, and an overall % score (above). However, this requires data to be entered electronically during or after audit.  
**Manual** - Count up the number of "Yes" answers and divide by the number of questions answered for that standard (excluding the "N/A"s), then multiply by 100 to get the percentage. To get the score for all the standards and for the audit overall, add the total number for each standard and divide by the number of questions and multiply by 100 to get the overall percentage.

\*developed using Infection Prevention Society (IPS) audit tools for monitoring IC guidelines within the community setting (2005); IPS self assessment audit for assessing implementation of HTM 01-05: decontamination in primary care dental practices and related infection prevention and control issues (2009); IPS audit tools for monitoring infection control standards (2004); International Scientific Forum on Home Hygiene (IFH) and IPS home hygiene - prevention of infection at home: a training resource for carers and their trainers (2003)

 [Click to download  
Word document](#)

## Literature review on the epidemiology of tattooing and its complications

V0.7

*Dr Victor Aiyedun*  
*Specialist Registrar,*  
*Public Health Medicine*

*Supervised by*  
*Dr Fortune Ncube*  
*Consultant Epidemiologist,*

Health Protection Agency, Colindale

1